

# **Introduction to Generalized Complex Geometry**



Publicações Matemáticas

**Introduction to Generalized  
Complex Geometry**

Gil Cavalcanti  
University of Oxford



26<sup>o</sup> Colóquio Brasileiro de Matemática

Copyright © 2007 by Gil R. Cavalcanti  
Direitos reservados, 2007 pela Associação Instituto  
Nacional de Matemática Pura e Aplicada - IMPA  
Estrada Dona Castorina, 110  
22460-320 Rio de Janeiro, RJ  
Impresso no Brasil / Printed in Brazil  
Capa: Noni Geiger / Sérgio R. Vaz

## **26<sup>a</sup> Colóquio Brasileiro de Matemática**

- Aspectos Ergódicos da Teoria dos Números - Alexander Arbieto, Carlos Matheus e Carlos Gustavo Moreira
- Componentes Irredutíveis dos Espaços de Folheações - Alcides Lins Neto
- Elliptic Regularity and Free Boundary Problems: an Introduction - Eduardo V. Teixeira
- Hiperbolicidade, Estabilidade e Caos em Dimensão Um - Flavio Abdenur e Luiz Felipe Nobili França
- **Introduction to Generalized Complex Geometry - Gil R. Cavalcanti**
- Introduction to Tropical Geometry - Grigory Mikhalkin
- Introdução aos Algoritmos Randomizados - Celina de Figueiredo, Guilherme da Fonseca, Manoel Lemos e Vinicius de Sá
- Mathematical Aspects of Quantum Field Theory - Edson de Faria and Wellington de Melo
- Métodos Estatísticos Não-Paramétricos e suas Aplicações - Aluisio Pinheiro e Hildete P. Pinheiro
- Moduli Spaces of Curves - Enrico Arbarello
- Noções de Informação Quântica - Marcelo O. Terra Cunha
- Three Dimensional Flows - Vítor Araújo e Maria José Pacifico
- Tópicos de Corpos Finitos com Aplicações em Criptografia e Teoria de Códigos - Ariane Masuda e Daniel Panario
- Tópicos Introdutórios à Análise Complexa Aplicada - André Nachbin e Ailín Ruiz de Zárate
- Uma Introdução à Mecânica Celeste - Sérgio B. Volchan
- Uma Introdução à Teoria Econômica dos Jogos - Humberto Bortolossi, Gilmar Garbugio e Brígida Sartini
- Uma Introdução aos Sistemas Dinâmicos via Frações Contínuas - Lorenzo J. Díaz e Danielle de Rezende Jorge

ISBN: 978-85-244-0254-8

**Distribuição:** IMPA  
Estrada Dona Castorina, 110  
22460-320 Rio de Janeiro, RJ  
E-mail: [ddic@impa.br](mailto:ddic@impa.br)  
<http://www.impa.br>

# Contents

<b>1</b>	<b>Generalized complex geometry</b>	<b>3</b>
1.1	Linear algebra . . . . .	5
1.1.1	Mukai pairing and pure forms . . . . .	7
1.1.2	The decomposition of forms . . . . .	9
1.1.3	The actions of $\mathcal{J}$ on forms . . . . .	10
1.2	The Courant bracket . . . . .	13
1.3	Generalized complex structures . . . . .	16
1.3.1	The differential graded algebra $(\Omega^\bullet(\bar{L}), d_L)$ . . . . .	20
1.4	Deformations . . . . .	21
1.5	Submanifolds . . . . .	24
1.6	Examples . . . . .	27
1.6.1	Symplectic fibrations . . . . .	28
1.6.2	Nilpotent Lie algebras . . . . .	30
1.6.3	A surgery . . . . .	30
<b>2</b>	<b>Generalized metric structures</b>	<b>33</b>
2.1	Linear algebra of the metric . . . . .	34
2.1.1	Generalized metric . . . . .	34
2.1.2	Hermitian structures . . . . .	36
2.2	Generalized Kähler structures . . . . .	38
2.2.1	Hermitian structures . . . . .	38
2.2.2	Generalized Kähler structures . . . . .	39
2.3	Hodge identities . . . . .	42
2.4	Formality . . . . .	43

<b>3</b>	<b>Reduction</b>	<b>47</b>
3.1	Courant algebras and extended actions . . . . .	49
3.1.1	Symmetries of Courant algebroids . . . . .	49
3.1.2	Extended actions . . . . .	50
3.1.3	Moment maps . . . . .	54
3.2	Reduction of Courant algebroids . . . . .	57
3.2.1	Examples . . . . .	58
3.3	Reduction of Dirac structures . . . . .	63
3.4	Reduction of gen. complex structures . . . . .	64
3.4.1	Symplectic structures . . . . .	67
3.4.2	Complex structures . . . . .	68
<b>4</b>	<b>T-duality</b>	<b>70</b>
4.1	T-duality with NS-flux . . . . .	72
4.1.1	Principal torus bundles . . . . .	74
4.2	T-duality as a map of Courant algebroids . . . . .	75
4.2.1	The metric and the Buscher rules . . . . .	79
4.2.2	The bihermitian structure . . . . .	80
4.3	Reduction and T-duality . . . . .	82
	<b>Bibliography</b>	<b>85</b>

# Chapter 1

## Generalized complex geometry

Generalized complex structures were introduced by Nigel Hitchin [37] and further developed by Gualtieri [34]. They are a simultaneous generalization of complex and symplectic structures obtained by searching for complex structures on  $TM \oplus T^*M$ , the sum of tangent and cotangent bundles of a manifold  $M$  or, more generally, on Courant algebroids over  $M$ .

Not only do generalized complex structures generalize symplectic and complex structures but also provide a unifying language for many features of these two seemingly distinct geometries. For instance, the operators  $\partial$  and  $\bar{\partial}$  and the  $(p, q)$ -decomposition of forms from complex geometry have their analogue in the generalized complex world as well as symplectic and Lagrangian submanifolds from the symplectic world.

This unifying property of generalized complex structures was immediately noticed by the physicists and the most immediate application was to mirror symmetry. From the generalized complex point of view, mirror symmetry should not be seen as the interchange of two different structures (complex to symplectic and vice versa) but just a transformation of the generalized complex structures in consideration. Features of mirror symmetry from the generalized complex

viewpoint were studied in [27, 33, 36] and in [4, 5, 18] from a more mathematical angle.

The relevance of generalized complex structures to string theory does not stop there. They also arise as solutions to the vacuum equations for some string theories, examples of which were given by Lindström, Minasian, Tomasiello and Zabzine [49] and Zucchini [69]. Furthermore, the generalized complex version of Kähler manifolds correspond to the bihermitian structures of Gates, Hull and Roček [30] obtained from the study of general  $(2, 2)$  supersymmetric sigma models (see also [50]).

Another angle to generalized complex structures comes from the study of Dirac structures: maximal isotropic subspaces  $L \subset TM \oplus T^*M$  together with an integrability condition. A generalized complex structure is nothing but a complex Dirac  $L \subset T_{\mathbb{C}}M \oplus T_{\mathbb{C}}^*M$  satisfying  $L \cap \bar{L} = \{0\}$ . Dirac structures predate generalized complex structure by more than 20 years and due to work of Weinstein and many of his collaborators we know an awful lot about them. Many of the features of generalized complex structures are in a way results about Dirac structures, e.g., some aspects of their local structure, the  $d_L$ -cohomology, the deformation theory and reduction procedure. However due to lack of space, we will not stress the connection between generalized complex geometry and Dirac structures.

This chapter follows closely the exposition of Gualtieri's thesis [34] and includes some developments to the theory obtained thereafter. This chapter is organized as follows. In the first section we introduce linear generalized complex structures, i.e., generalized complex structures on vector spaces and then go on to show that these structures give rise to a decomposition of forms similar to the  $(p, q)$ -decomposition of forms in a complex manifold. In Section 1.2, we introduce the Courant bracket which furnishes the integrability condition for a generalized complex structure on a manifold, as we see in Section 1.3. This is a compatibility condition between the pointwise defined generalized complex structure and the differential structure, which is equivalent to saying that the pointwise decomposition of forms induced by the generalized complex structure gives rise to a decomposition of the exterior derivative  $d = \partial + \bar{\partial}$ . In Section 1.4 we state the basic result on the deformation theory of generalized complex structures and in Section 1.5 we study two important classes of



submanifolds of a generalized complex manifold. We finish studying some interesting examples of generalized complex manifolds in the last section.

## 1.1 Linear algebra of a generalized complex structure

For any vector space  $V^n$  we define *the double of  $V$* ,  $\mathcal{D}V$ , to be a  $2n$ -dimensional vector space endowed with a nondegenerate pairing  $\langle \cdot, \cdot \rangle$  and a surjective projection

$$\pi : \mathcal{D}V \longrightarrow V,$$

such that the kernel of  $\pi$  is isotropic. Observe that the requirement that the kernel of  $\pi$  is isotropic implies that the pairing has signature  $(n, n)$ .

Using the pairing to identify  $(\mathcal{D}V)^*$  with  $\mathcal{D}V$ , we get a map

$$\frac{1}{2}\pi^* : V^* \longrightarrow \mathcal{D}V,$$

so we can regard  $V^*$  as a subspace of  $\mathcal{D}V$ . By definition,  $\langle \pi^*(V^*), \text{Ker}(\pi) \rangle = 0$  and, since  $\text{Ker}(\pi)^\perp = \text{Ker}(\pi)$ , we see that  $\pi^*(V^*) = \text{Ker}(\pi)$ , therefore furnishing the following exact sequence

$$0 \longrightarrow V^* \xrightarrow{\frac{1}{2}\pi^*} \mathcal{D}V \xrightarrow{\pi} V \longrightarrow 0.$$

If we choose an isotropic splitting  $\nabla : V \longrightarrow \mathcal{D}V$ , i.e., a splitting for which  $\nabla(V)$  is isotropic, then we obtain an isomorphism  $\mathcal{D}V \cong V \oplus V^*$  and the pairing is nothing but the natural pairing on  $V \oplus V^*$ :

$$\langle X + \xi, Y + \eta \rangle = \frac{1}{2}(\xi(Y) + \eta(X)).$$

**Definition 1.1.** A *generalized complex structure* on  $V$  is a linear complex structure  $\mathcal{J}$  on  $\mathcal{D}V$  orthogonal with respect to the pairing.

Since  $\mathcal{J}^2 = -\text{Id}$ , it splits  $\mathcal{D}V \otimes \mathbb{C}$  as a direct sum of  $\pm i$ -eigenspaces,  $L$  and  $\bar{L}$ . Further, as  $\mathcal{J}$  is orthogonal, we obtain that for  $v, w \in L$ ,

$$\langle v, w \rangle = \langle \mathcal{J}v, \mathcal{J}w \rangle = \langle iv, iw \rangle = -\langle v, w \rangle,$$

and hence  $L$  is a maximal isotropic subspace with respect to the pairing.

Conversely, prescribing such an  $L$  as the  $i$ -eigenspace determines a unique generalized complex structure on  $V$ , therefore a generalized complex structure on a vector space  $V^n$  is equivalent to a maximal isotropic subspace  $L \subset \mathcal{D}V \otimes \mathbb{C}$  such that  $L \cap \bar{L} = \{0\}$

This last point of view also shows that a generalized complex structure is a special case of a more general object called a *Dirac structure*, which is a maximal isotropic subspace of  $\mathcal{D}V$ . So a generalized complex structure is nothing but a complex Dirac structure  $L$  for which  $L \cap \bar{L} = \{0\}$ .

**Example 1.2** (Complex structures). If we have a splitting  $\mathcal{D}V = V \oplus V^*$  and  $V$  has a complex structure  $I$ , then it induces a generalized complex structure on  $V$  which can be written in matrix form using the splitting as

$$\mathcal{J}_I = \begin{pmatrix} -I & 0 \\ 0 & I^* \end{pmatrix}.$$

The  $i$ -eigenspace of  $\mathcal{J}_I$  is  $L = V^{0,1} \oplus V^{*,1,0} \subset (V \oplus V^*) \otimes \mathbb{C}$ . It is clear that  $L$  is a maximal isotropic subspace and that  $L \cap \bar{L} = \{0\}$ .

**Example 1.3** (Symplectic structures). Again, if we have a splitting  $\mathcal{D}V = V \oplus V^*$ , then a symplectic structure  $\omega$  on  $V$  also induces a generalized complex structure  $\mathcal{J}_\omega$  on  $V$  by letting

$$\mathcal{J}_\omega = \begin{pmatrix} 0 & -\omega^{-1} \\ \omega & 0 \end{pmatrix}.$$

The  $i$ -eigenspace of  $\mathcal{J}_\omega$  is given by  $L = \{X - i\omega(X) : X \in V\}$ . The nondegeneracy of  $\omega$  implies that  $L \cap \bar{L} = \{0\}$  and skew symmetry implies that  $L$  is isotropic.

**Example 1.4.** A real 2-form  $B$  acts naturally on  $\mathcal{D}V$  by the *B-field transform*

$$e \mapsto e - B(\pi(e)).$$

If  $V$  is endowed with a generalized complex structure,  $\mathcal{J}$ , whose  $+i$ -eigenspace is  $L$ , we can consider its image under the action of a  $B$ -field:  $L_B = \{e - B(\pi(e)) : e \in L\}$ . Since  $B$  is real,  $L_B \cap \bar{L}_B = (\text{Id} - B)L \cap \bar{L} = \{0\}$ . Again, skew symmetry implies that  $L_B$  is

isotropic. If we have a splitting for  $\mathcal{D}V$ , we can write  $\mathcal{J}_B$ , the  $B$ -field transform of  $\mathcal{J}$ , in matrix form

$$\mathcal{J}_B = \begin{pmatrix} 1 & 0 \\ -B & 1 \end{pmatrix} \mathcal{J} \begin{pmatrix} 1 & 0 \\ B & 1 \end{pmatrix}.$$

One can check that any two isotropic splittings of  $\mathcal{D}V$  are related by  $B$ -field transforms.

**Example 1.5.** In the presence of a splitting  $\mathcal{D}V = V \oplus V^*$ , an element  $\beta \in \wedge^2 V$  also acts in a similar fashion:

$$X + \xi \mapsto X + \xi + \xi \lrcorner \beta.$$

An argument similar to the one above shows that the  $\beta$ -transform of a generalized complex structure is still a generalized complex structure.

### 1.1.1 Mukai pairing and pure forms

In the presence of a splitting  $\mathcal{D}V = V \oplus V^*$ , we have one more characterization of a generalized complex structure on  $V$ , obtained from an interpretation of forms as spinors.

The Clifford algebra of  $\mathcal{D}V$  is defined using the natural form  $\langle \cdot, \cdot \rangle$ , i.e., for  $v \in \mathcal{D}V \subset \text{Cl}(\mathcal{D}V)$  we have  $v^2 = \langle v, v \rangle$ . Since  $V^*$  is a maximal isotropic, its exterior algebra is a subalgebra of  $\text{Cl}(\mathcal{D}V)$ . In particular,  $\wedge^n V^*$  is a distinguished line in the Clifford algebra and generates a left ideal  $\mathcal{I}$ . A splitting  $\mathcal{D}V = V \oplus V^*$  gives an isomorphism  $\mathcal{I} \cong \wedge^\bullet V \cdot \wedge^n V^* \cong \wedge^\bullet V^*$ . This, in turn, determines an action of the Clifford algebra on  $\wedge^\bullet V^*$  by

$$(X + \xi) \cdot \alpha = i_X \alpha + \xi \wedge \alpha.$$

If we let  $\sigma$  be the antiautomorphism of  $\text{Cl}(V \oplus V^*)$  defined on decomposables by

$$\sigma(v_1 \cdot v_2 \cdots v_k) = v_k \cdots v_2 \cdot v_1, \quad (1.1)$$

then we have the following bilinear form on  $\wedge^\bullet V^* \subset \text{Cl}(V \oplus V^*)$ :

$$(\xi_1, \xi_2) \mapsto (\sigma(\xi_1) \wedge \xi_2)_{\text{top}},$$

where *top* indicates taking the top degree component on the form. If we decompose  $\xi_i$  by degree:  $\xi_i = \sum \xi_i^j$ , with  $\deg(\xi_i^j) = j$ , the above can be rewritten, in an  $n$ -dimensional space, as

$$(\xi_1, \xi_2) = \sum_j (-1)^j (\xi_1^{2j} \wedge \xi_2^{n-2j} + \xi_1^{2j+1} \wedge \xi_2^{n-2j-1}) \quad (1.2)$$

This bilinear form coincides in cohomology with the *Mukai pairing*, introduced in a  $K$ -theoretical framework in [56].

Now, given a form  $\rho \in \wedge^\bullet V^* \otimes \mathbb{C}$  (of possibly mixed degree) we can consider its Clifford annihilator

$$L_\rho = \{v \in (V \oplus V^*) \otimes \mathbb{C} : v \cdot \rho = 0\}.$$

It is clear that  $\overline{L_\rho} = L_{\bar{\rho}}$ . Also, for  $v \in L_\rho$ ,

$$0 = v^2 \cdot \rho = \langle v, v \rangle \rho,$$

thus  $L_\rho$  is always isotropic.

**Definition 1.6.** An element  $\rho \in \wedge^\bullet V^*$  is a *pure form* if  $L_\rho$  is maximal, i.e.,  $\dim_{\mathbb{C}} L_\rho = \dim_{\mathbb{R}} V$ .

Given a maximal isotropic subspace  $L \subset V \oplus V^*$  one can always find a pure form annihilating it and conversely, if two pure forms annihilate the same maximal isotropic, then they are a multiple of each other, i.e., maximal isotropics are in one-to-one correspondence with lines of pure forms. Algebraically, the requirement that a form is pure implies that it is of the form  $e^{B+i\omega}\Omega$ , where  $B$  and  $\omega$  are real 2-forms and  $\Omega$  is a decomposable complex form. The relation between the Mukai pairing and generalized complex structures comes in the following:

**Proposition 1.7.** (Chevalley [21]) *Let  $\rho$  and  $\tau$  be pure forms. Then  $L_\rho \cap L_\tau = \{0\}$  if and only if  $(\rho, \tau) \neq 0$ .*

Therefore a pure form  $\rho = e^{B+i\omega}\Omega$  determines a generalized complex structure if and only if  $(\rho, \bar{\rho}) \neq 0$ . This is only possible if  $V$  is even dimensional, say  $\dim(V) = 2n$ , in which case  $(\rho, \bar{\rho}) = \Omega \wedge \bar{\Omega} \wedge \omega^{n-k}$ , where  $k$  is the degree of  $\Omega$ . In particular, there is no generalized complex structure on odd dimensional spaces.

With this we see that, if  $\mathcal{D}V$  is split, then a generalized complex structure is equivalent to a line  $K \subset \wedge^\bullet V^* \otimes \mathbb{C}$  generated by a form  $e^{B+i\omega}\Omega$ , such that  $\Omega$  is a decomposable complex form of degree, say,  $k$ ,  $B$  and  $\omega$  are real 2-forms and  $\Omega \wedge \bar{\Omega} \wedge \omega^{n-k} \neq 0$ . The degree of the form  $\Omega$  is the *type* of the generalized complex structure. The line  $K$  annihilating  $L$  is the *canonical line*.

EXAMPLES 1.2 – 1.5 REVISED: The canonical line in  $\wedge^\bullet V^* \otimes \mathbb{C}$  that gives the generalized complex structure for a complex structure is  $\wedge^{n,0}V^*$ , while the line for a symplectic structure  $\omega$  is generated by  $e^{i\omega}$ . If  $\rho$  is a generator of the canonical line of a generalized complex structure  $\mathcal{J}$ ,  $e^B \wedge \rho$  is a generator of a  $B$ -field transform of  $\mathcal{J}$  and  $e^{\beta \lrcorner} \rho$  is a generator for a  $\beta$ -field transform of  $\mathcal{J}$ .

### 1.1.2 The decomposition of forms

Using the same argument used before with  $V$  and  $V^*$  to the maximal isotropics  $L$  and  $\bar{L}$  determining a generalized complex structure on  $V$ , we see that  $\text{Cl}((V \oplus V^*) \otimes \mathbb{C}) \cong \text{Cl}(L \oplus \bar{L})$  acts on  $\wedge^{2n}L$  and the left ideal generated is the subalgebra  $\wedge^\bullet \bar{L}$ . The choice of a pure form  $\rho$  for the generalized complex structure gives an isomorphism of Clifford modules:

$$\phi : \wedge^\bullet \bar{L} \longrightarrow \wedge^\bullet V^* \otimes \mathbb{C}; \quad \phi(s) = s \cdot \rho.$$

The decomposition of  $\wedge^\bullet \bar{L}$  by degree gives rise to a new decomposition of  $\wedge^\bullet V^* \otimes \mathbb{C}$  and the Mukai pairing on  $\wedge^\bullet V^* \otimes \mathbb{C}$  corresponds to the same pairing on  $\wedge^\bullet \bar{L}$ . But in  $\wedge^\bullet \bar{L}$  the Mukai pairing is nondegenerate in

$$\wedge^k \bar{L} \times \wedge^{2n-k} \bar{L} \longrightarrow \wedge^{2n} \bar{L}$$

and vanishes in  $\wedge^k \bar{L} \times \wedge^l \bar{L}$  for all other values of  $l$ . Therefore the same is true for the induced decomposition on forms.

**Proposition 1.8.** *Letting  $U^k = \wedge^{n-k} \bar{L} \cdot \rho \subset \wedge^\bullet V^* \otimes \mathbb{C}$ , the Mukai pairing on  $U^k \times U^l$  is trivial unless  $l = -k$ , in which case it is nondegenerate.*

According to the definition above,  $U^n$  is the canonical line. Also, the elements of  $U^k$  are even/odd forms, according to the parity of  $k$ ,

$n$  and the type of the structure. For example, if  $n$  and the type are even, the elements of  $U^k$  will be even if and only if  $k$  is even.

**Example 1.9.** In the complex case, we take  $\rho \in \wedge^{n,0}V^* \setminus \{0\}$  to be generator of the canonical line of the induced generalized complex structure. Then, from Example 1.2, we have that  $\bar{L} = \wedge^{1,0}V \oplus \wedge^{0,1}V^*$ , so

$$U^k = \oplus_{p-q=k} \wedge^{p,q} V^*.$$

Then, in this case, one can see Proposition 1.8 as a consequence of the fact that the top degree part of the exterior product vanishes on  $\wedge^{p,q}V^* \times \wedge^{p',q'}V^*$ , unless  $p + p' = q + q' = n$ , in which case it is a nondegenerate pairing.

**Example 1.10.** The decomposition of forms into the spaces  $U^k$  for a symplectic vector space  $(V, \omega)$  was worked out in [16]. In this case we have,

$$U^k = \{e^{i\omega} e^{\frac{\Lambda}{2i}} \wedge^{n-k} V^* \otimes \mathbb{C}\},$$

where  $\Lambda$  is the Poisson bivector associated to the symplectic structure,  $\Lambda = -\omega^{-1}$ , and acts on forms by interior product.

So  $\Phi : \wedge^{n-k}V^* \otimes \mathbb{C} \longrightarrow \wedge^{n-k}V^* \otimes \mathbb{C}$  defined by

$$\Phi(\alpha) = e^{i\omega} e^{\frac{\Lambda}{2i}} \alpha$$

is a natural isomorphism of graded spaces:  $\Phi(\wedge^k V^* \otimes \mathbb{C}) = U^{n-k}$ .

**Example 1.11.** If a generalized complex structure induces a decomposition of the differential forms into the spaces  $U^k$ , then the  $B$ -field transform of this structure will induce a decomposition into  $U_B^k = e^B \wedge U^k$ . Indeed, by Example 1.4 revised,  $U_B^n = e^B \wedge U^n$ , and

$$U_B^k = (\text{Id} - B)(\bar{L}) \cdot U_B^{k+1}.$$

The desired expression can be obtained by induction.

### 1.1.3 The actions of $\mathcal{J}$ on forms

Recall that the group  $\text{Spin}(n, n)$  sits inside  $\text{Cl}(V \oplus V^*)$  as

$$\text{Spin}(n, n) = \{v_1 \cdots v_{2k} : v_i \cdot v_i = \pm 1; k \in \mathbb{N}\}.$$

And  $\text{Spin}(n, n)$  is a double cover of  $\text{SO}(n, n)$ :

$$\varphi : \text{Spin}(n, n) \longrightarrow \text{SO}(n, n); \quad \varphi(v)X = v \cdot X \cdot \sigma(v),$$

where  $\sigma$  is the main antiautomorphism of the Clifford algebra as defined in (1.1).

This map identifies the Lie algebras  $\mathfrak{spin}(n, n) \cong \mathfrak{so}(n, n) \cong \wedge^2 V \oplus \wedge^2 V^* \oplus \text{End}(V)$ :

$$\mathfrak{spin}(n, n) \longrightarrow \mathfrak{so}(n, n); \quad d\varphi(v)(X) = [v, X] = v \cdot X - X \cdot v$$

But, as the exterior algebra of  $V^*$  is naturally the space of spinors, each element in  $\mathfrak{spin}(n, n)$  acts naturally on  $\wedge^\bullet V^*$ .

**Example 1.12.** Let  $B = \sum b_{ij}e^i \wedge e^j \in \wedge^2 V^* \subset \mathfrak{so}(n, n)$  be a 2-form. As an element of  $\mathfrak{so}(n, n)$ ,  $B$  acts on  $V \oplus V^*$  via

$$X + \xi \mapsto X \lrcorner B.$$

Then the corresponding element in  $\mathfrak{spin}(n, n)$  inducing the same action on  $V \oplus V^*$  is given by  $\sum b_{ij}e^j e^i$ , since, in  $\mathfrak{so}(n, n)$ , we have

$$e^i \wedge e^j : e_k \mapsto \delta_{ik}e^j - \delta_{jk}e^i.$$

And, in  $\mathfrak{spin}(n, n)$ ,

$$d\varphi(e^j e^i)e_k = (e^j e^i) \cdot e_k - e_k \cdot (e^j e^i) = e^j \cdot (e^i \cdot e_k) - (e_k \cdot e^j) \cdot e^i = \delta_{ik}e^j - \delta_{jk}e^i.$$

Finally, the spinorial action of  $B$  on a form  $\varphi$  is given by

$$\sum b_{ij}e^j e^i \cdot \varphi = -B \wedge \varphi.$$

**Example 1.13.** Similarly, for  $\beta = \sum \beta^{ij}e_i \wedge e_j \in \wedge^2 V \subset \mathfrak{so}(n, n)$ , its action is given by

$$\beta \cdot (X + \xi) = i_\xi \beta.$$

And the corresponding element in  $\mathfrak{spin}(n, n)$  with the same action is  $\sum \beta^{ij}e_j e_i$ . The action of this element on a form  $\varphi$  is given by

$$\beta \cdot \varphi = \beta \lrcorner \varphi.$$

**Example 1.14.** Finally, an element of  $A = \sum A_i^j e^i \otimes e_j \in \text{End}(V) \subset \mathfrak{so}(n, n)$  acts on  $V \oplus V^*$  via

$$A(X + \xi) = A(X) + A^*(\xi).$$

The element of  $\mathfrak{spin}(n, n)$  with the same action is  $\frac{1}{2} \sum A_i^j (e_j e^i - e^i e_j)$ . And the action of this element on a form  $\varphi$  is given by:

$$\begin{aligned} A \cdot \varphi &= \frac{1}{2} \sum A_i^j (e_j e^i - e^i e_j) \cdot \varphi \\ &= \frac{1}{2} \sum A_i^j (e_j \lrcorner (e^i \wedge \varphi) - e^i \wedge (e_j \lrcorner \varphi)) \\ &= \frac{1}{2} \sum_i A_i^i \varphi - \sum_{i,j} A_i^j e^i \wedge (e_j \lrcorner \varphi) \\ &= -A^* \varphi + \frac{1}{2} \text{Tr} A \varphi, \end{aligned}$$

where  $A^* \varphi$  is the Lie algebra adjoint of  $A$  action of  $\varphi$  via

$$A^* \varphi(v_1, \dots, v_p) = \sum_i \varphi(v_1, \dots, A v_i, \dots, v_p).$$

The reason for introducing this Lie algebra action of  $\mathfrak{spin}(n, n)$  on forms is because  $\mathcal{J} \in \mathfrak{spin}(n, n)$ , hence we can compute its action on forms.

**Example 1.15.** In the case of a generalized complex structure induced by a symplectic one, we have that  $\mathcal{J}$  is the sum of a 2-form,  $\omega$ , and a bivector,  $-\omega^{-1}$ , hence its Lie algebra action on a form  $\varphi$  is

$$\mathcal{J} \varphi = (-\omega \wedge -\omega^{-1} \lrcorner) \varphi \quad (1.3)$$

**Example 1.16.** If  $\mathcal{J}$  is a generalized complex structure on  $V$  induced by a complex structure  $I$ , then its Lie algebra action is the one corresponding to the traceless endomorphism  $-I$  (see Example 1.2). Therefore, when acting on a  $(p, q)$ -form  $\alpha$ :

$$\mathcal{J} \cdot \alpha = I^* \alpha = i(p - q) \alpha.$$

From this example and Example 1.9 it is clear that in the case of a generalized complex structure induced by a complex one, the subspaces  $U^k \subset \wedge^\bullet V^*$  are the  $ik$ -eigenspaces of the action of  $\mathcal{J}$ . This is general.



**Proposition 1.17.** *The spaces  $U^k$  are the  $ik$ -eigenspaces of the Lie algebra action of  $\mathcal{J}$ .*

*Proof.* Recall from Subsection 1.1.2 that the choice of a nonzero element  $\rho$  of the canonical line  $K \subset \wedge^\bullet V^*$  gives an isomorphism of Clifford modules:

$$\phi : \wedge^\bullet \bar{L} \longrightarrow \wedge^\bullet V^* \otimes \mathbb{C}; \quad \phi(s) = s \cdot \rho.$$

And the spaces  $U^k$  are defined as  $U^{n-k} = \phi(\wedge^k \bar{L})$ . Further,  $\mathcal{J}$  acts on  $L^* \cong \bar{L}$  as multiplication by  $-i$ . Hence, by Example 1.14, its Lie algebra action on  $\gamma \in \wedge^k \bar{L}$  is:

$$\mathcal{J} \cdot \gamma = -J^* \gamma + \frac{1}{2} \text{Tr} J \gamma = -ik\gamma + \frac{1}{2} 2in\gamma = i(n-k)\gamma.$$

As  $\phi$  is an isomorphism of Clifford modules, for  $\alpha \in U^{n-k}$  we have  $\phi^{-1}\alpha \in \wedge^k \bar{L}$  and

$$\mathcal{J} \cdot \alpha = \mathcal{J} \cdot \phi(\phi^{-1}\alpha) = \phi(\mathcal{J}\phi^{-1} \cdot \alpha) = i(n-k)\phi(\phi^{-1}\alpha) = i(n-k)\alpha.$$

□

## 1.2 The Courant bracket and Courant algebroids

From the linear algebra developed in the previous section, it is clear that a generalized complex structure on a manifold lives naturally on the double of the tangent bundle,  $\mathcal{DT}$ . Similarly to the case of complex structures on a manifold, the integrability condition for a generalized complex structure is that its  $i$ -eigenspace has to be closed under a certain bracket known as the *Courant bracket*, originally introduced by Courant and Weinstein as an extension of the Lie bracket of vector fields to sections of  $TM \oplus T^*M$  [23, 22].

One of the striking features of the Courant bracket is that it only satisfies the Jacobi identity modulo an exact element, more precisely, for all  $e_1, e_2, e_3 \in \Gamma(TM \oplus T^*M)$  we have

$$\text{Jac}(e_1, e_2, e_3) := \llbracket \llbracket e_1, e_2 \rrbracket, e_3 \rrbracket + c.p. = \frac{1}{3} d(\llbracket \llbracket e_1, e_2 \rrbracket, e_3 \rrbracket + c.p.).$$

Liu, Weinstein and Xu, [51], axiomatized the properties of the Courant bracket in the concept of a Courant algebroid, which is the central object of this section. As we will see later, *exact* Courant algebroids are the natural space where generalized complex structures live.

**Definition 1.18.** A *Courant algebroid* over a manifold  $M$  is a vector bundle  $\mathcal{E} \rightarrow M$  equipped with a skew-symmetric bracket  $\llbracket \cdot, \cdot \rrbracket$  on  $\Gamma(\mathcal{E})$ , a nondegenerate symmetric bilinear form  $\langle \cdot, \cdot \rangle$ , and a bundle map  $\pi : \mathcal{E} \rightarrow TM$ , which satisfy the following conditions for all  $e_1, e_2, e_3 \in \Gamma(\mathcal{E})$  and  $f, g \in C^\infty(M)$ :

- C1)  $\pi(\llbracket e_1, e_2 \rrbracket) = [\pi(e_1), \pi(e_2)],$
- C2)  $\text{Jac}(e_1, e_2, e_3) = \frac{1}{3}d(\langle \llbracket e_1, e_2 \rrbracket, e_3 \rangle + c.p.),$
- C3)  $\llbracket e_1, f e_2 \rrbracket = f \llbracket e_1, e_2 \rrbracket + (\pi(e_1)f)e_2 - \langle e_1, e_2 \rangle df,$
- C4)  $\pi \circ d = 0$ , i.e.  $\langle df, dg \rangle = 0,$
- C5)  $\pi(e_1)\langle e_2, e_3 \rangle = \langle e_1 \bullet e_2, e_3 \rangle + \langle e_2, e_1 \bullet e_3 \rangle,$

where we consider  $\Omega^1(M)$  as a subset of  $\Gamma(\mathcal{E})$  via the map  $\frac{1}{2}\pi^* : \Omega^1(M) \rightarrow \Gamma(\mathcal{E})$  (using  $\langle \cdot, \cdot \rangle$  to identify  $\mathcal{E}$  with  $\mathcal{E}^*$ ) and  $\bullet$  denotes the combination

$$e_1 \bullet e_2 = \llbracket e_1, e_2 \rrbracket + d\langle e_1, e_2 \rangle. \quad (1.4)$$

and is the *adjoint action* of  $e_1$  on  $e_2$ .

**Definition 1.19.** A Courant algebroid is *exact* if the following sequence is exact:

$$0 \longrightarrow T^*M \xrightarrow{\frac{1}{2}\pi^*} \mathcal{E} \xrightarrow{\pi} TM \longrightarrow 0 \quad (1.5)$$

Given an exact Courant algebroid, we may always choose an isotropic right splitting  $\nabla : TM \rightarrow \mathcal{E}$ . Such a splitting has a *curvature* 3-form  $H \in \Omega_{cl}^3(M)$  defined as follows, for  $X, Y \in \Gamma(TM)$ :

$$H(X, Y, Z) = \frac{1}{2}\langle \llbracket \nabla(X), \nabla(Y) \rrbracket, \nabla(Z) \rangle. \quad (1.6)$$

Using the bundle isomorphism  $\nabla + \frac{1}{2}\pi^* : TM \oplus T^*M \rightarrow \mathcal{E}$ , we transport the Courant algebroid structure onto  $TM \oplus T^*M$ . As before the pairing is nothing but the natural pairing

$$\langle X + \xi, Y + \eta \rangle = \frac{1}{2}(\eta(X) + \xi(Y)), \quad (1.7)$$

and for  $X + \xi, Y + \eta \in \Gamma(TM \oplus T^*M)$  the bracket becomes

$$[[X + \xi, Y + \eta]] = [X, Y] + \mathcal{L}_X \eta - \mathcal{L}_Y \xi - \frac{1}{2}d(i_X \eta - i_Y \xi) + i_Y i_X H, \quad (1.8)$$

which is the *H-twisted Courant bracket* on  $TM \oplus T^*M$  [59]. Isotropic splittings of (1.5) differ by 2-forms  $b \in \Omega^2(M)$ , and a change of splitting modifies the curvature  $H$  by the exact form  $db$ . Hence the cohomology class  $[H] \in H^3(M, \mathbb{R})$ , called the *characteristic class* or *Ševera class* of  $\mathcal{E}$ , is independent of the splitting and determines the exact Courant algebroid structure on  $\mathcal{E}$  completely.

One way to see the Courant bracket on  $TM \oplus T^*M$  as a natural extension of the Lie bracket of vector fields is as follows. Recall that the Lie bracket satisfies (and can be defined by) the following identity when acting on a form  $\alpha$  (see [44], Chapter 1, Proposition 3.10):

$$2i_{[v_1, v_2]} \alpha = i_{v_1 \wedge v_2} d\alpha + d(i_{v_1 \wedge v_2} \alpha) + 2i_{v_1} d(i_{v_2} \alpha) - 2i_{v_2} d(i_{v_1} \alpha).$$

Now we observe that this formula gives a natural extension of the Lie bracket to a bracket on  $TM \oplus T^*M$ , acting on forms via the Clifford action:

$$2[[v_1, v_2] \cdot \alpha] = v_1 \wedge v_2 \cdot d\alpha + d(v_1 \wedge v_2 \cdot \alpha) + 2v_1 \cdot d(v_2 \cdot \alpha) - 2v_2 \cdot d(v_1 \cdot \alpha). \quad (1.9)$$

Spelling it out we obtain (see Gualtieri [34], Lemma 4.24):

$$[[X + \xi, Y + \eta]] = [X, Y] + \mathcal{L}_X \eta - \mathcal{L}_Y \xi - \frac{1}{2}d(i_X \eta - i_Y \xi).$$

This is the Courant bracket with  $H = 0$ . If we replace  $d$  by  $d_H = d + H \wedge$  in (1.9) we obtain the *H-twisted Courant bracket*:

$$2[[v_1, v_2]_H \cdot \alpha] = v_1 \wedge v_2 \cdot d_H \alpha + d_H(v_1 \wedge v_2 \cdot \alpha) + 2v_1 \cdot d_H(v_2 \cdot \alpha) - 2v_2 \cdot d_H(v_1 \cdot \alpha). \quad (1.10)$$

### 1.3 Generalized complex structures

Given the work in the previous sections it is clear that the fiber of an exact Courant algebroid  $\mathcal{E}$  over a point  $p \in M$  is nothing but  $DT_pM$  and hence it is natural to define a *generalized almost complex structure* as a differentiable bundle automorphism  $\mathcal{J} : \mathcal{E} \rightarrow \mathcal{E}$  which is a linear generalized complex structure on each fiber. The Courant bracket provides the integrability condition.

**Definition 1.20.** A *generalized complex structure* on an exact Courant algebroid  $\mathcal{E}$  is a generalized almost complex structure  $\mathcal{J}$  on  $\mathcal{E}$  whose  $i$ -eigenspace is closed with respect to the Courant bracket.

As before,  $\mathcal{J}$  can be described in terms of its  $i$ -eigenspace,  $L$ , which is a maximal isotropic subspace of  $\mathcal{E}_{\mathbb{C}}$  closed under the Courant bracket satisfying  $L \cap \bar{L} = \{0\}$ .

The choice of a splitting for  $\mathcal{E}$  making it isomorphic to  $TM \oplus T^*M$  with the  $H$ -Courant bracket also allows us to characterize a generalized complex structure in terms of its *canonical bundle*,  $K \subset \wedge^{\bullet} T_{\mathbb{C}}^*M$ , the line subbundle of  $\wedge T_{\mathbb{C}}^*M$  whose fiber over  $p$  is the canonical line for the generalized complex structure on  $\mathcal{E}_p \cong DT_pM$ . If  $\rho$  is a non-vanishing local section of  $K$  then using (1.10) one can easily see that the integrability condition is equivalent to the existence of a local section  $e \in \Gamma(T_{\mathbb{C}}M \oplus T_{\mathbb{C}}^*M)$  such that

$$d_H \rho = e \cdot \rho.$$

If we let  $\mathcal{U}^k$  be the space of sections of the bundle  $U^k$ , defined in Proposition 1.8, this is only the case if  $d_H \rho \in \mathcal{U}^{n-1} = \bar{L} \cdot \mathcal{U}^n$ .

**Example 1.21.** An almost complex structure on a manifold  $M$  induces a generalized almost complex structure with  $i$ -eigenspace  $T^{0,1}M \oplus T^{*1,0}M$ . If this generalized almost complex structure is integrable, then  $T^{0,1}M$  has to be closed with respect to the Lie bracket and hence the almost complex structure is actually a complex structure. Conversely, any complex structure gives rise to an integrable generalized complex structure.

**Example 1.22.** If  $M$  has a nondegenerate 2-form  $\omega$ , then the induced generalized almost complex structure will be integrable if for

some  $X + \xi$  we have

$$de^{i\omega} = (X + \xi) \cdot e^{i\omega}.$$

The degree 1 part gives that  $X \lrcorner \omega + \xi = 0$  and the degree 3 part, that  $d\omega = 0$  and hence  $M$  is a symplectic manifold.

**Example 1.23.** The action of a real closed 2-form  $B$  by  $B$ -field transforms on an exact Courant algebroid  $\mathcal{E}$  is a symmetry of the bracket. In fact,  $B$ -field transforms together with diffeomorphisms of the manifold, form the group of orthogonal symmetries of the Courant bracket. Therefore we can always transform a given a generalized complex structure by  $B$ -fields to obtain a new generalized complex structure which should be considered equivalent to the first one.

Assume we have a splitting for  $\mathcal{E}$  rendering it isomorphic to  $TM \oplus T^*M$  with the  $H$ -Courant bracket. If the 2-form  $B$  is not closed, then it induces an isomorphism between the  $H$ -Courant bracket and the  $H + dB$ -Courant bracket. In particular, if  $[H] = 0 \in H^3(M, \mathbb{R})$ , the bracket  $[\cdot, \cdot]_H$  is isomorphic to  $[\cdot, \cdot]_0$  by the action of a nonclosed 2-form.

**Example 1.24.** Consider  $\mathbb{C}^2$  with complex coordinates  $z_1, z_2$ . The differential form

$$\rho = z_1 + dz_1 \wedge dz_2$$

is equal to  $dz_1 \wedge dz_2$  along the locus  $z_1 = 0$ , while away from this locus it can be written as

$$\rho = z_1 \exp\left(\frac{dz_1 \wedge dz_2}{z_1}\right). \quad (1.11)$$

Since it also satisfies  $d\rho = -\partial_2 \cdot \rho$ , we see that it generates a canonical bundle  $K$  for a generalized complex structure which has type 2 along  $z_1 = 0$  and type 0 elsewhere, showing that a generalized complex structure does not necessarily have constant type.

In order to obtain a compact type-change locus we observe that this structure is invariant under translations in the  $z_2$  direction, hence we can take a quotient by the standard  $\mathbb{Z}^2$  action to obtain a generalized complex structure on the torus fibration  $D^2 \times T^2$ , where  $D^2$  is the unit disc in the  $z_1$ -plane. Using polar coordinates,  $z_1 = re^{2\pi i\theta_1}$ , the canonical bundle is generated, away from the central fibre, by

$$\exp(B + i\omega) = \exp(d \log r + id\theta_1) \wedge (d\theta_2 + id\theta_3)$$

and hence

$$\begin{aligned} B &= d \log r \wedge d\theta_2 - d\theta_1 \wedge d\theta_3 \\ \omega &= d \log r \wedge d\theta_3 + d\theta_1 \wedge d\theta_2, \end{aligned} \tag{1.12}$$

where  $\theta_2$  and  $\theta_3$  are coordinates for the 2-torus with unit periods. Away from  $r = 0$ , therefore, the structure is a  $B$ -field transform of the symplectic structure  $\omega$ .

The type jumps from 0 to 2 along the central fibre  $r = 0$ , inducing a complex structure on the restricted tangent bundle, for which the tangent bundle to the fibre is a complex sub-bundle. Hence the type change locus inherits the structure of a smooth elliptic curve with Teichmüller parameter  $\tau = i$ .

Similarly to the complex case, the integrability condition places restrictions on  $d_H(\mathcal{U}^k)$  for every  $k$  and hence allows us to define operators  $\partial$  and  $\bar{\partial}$ .

**Theorem 1.25** (Gualtieri [34], Theorem 4.3). *A generalized almost complex structure is integrable if and only if  $d_H : \mathcal{U}^k \rightarrow \mathcal{U}^{k+1} \oplus \mathcal{U}^{k-1}$ .*

*Proof.* It is clear that if  $d_H : \mathcal{U}^k \rightarrow \mathcal{U}^{k+1} \oplus \mathcal{U}^{k-1}$  for all  $k$ , then, in particular, for  $k = n$  we get the integrability condition

$$d_H : \mathcal{U}^n \rightarrow \mathcal{U}^{n-1},$$

as  $\mathcal{U}^{n+1} = \{0\}$ .

In order to obtain the converse we shall first prove that

$$d_H : \mathcal{U}^k \rightarrow \oplus_{j \geq k-1} \mathcal{U}^j. \tag{1.13}$$

This is done by induction on  $k$ ,  $k$  starting at  $n$  and going downwards. The first step is just the integrability condition. Assuming that the claim is true for  $k' > k$ , let  $v_1, v_2$  be sections of  $L$  and  $\varphi \in \mathcal{U}^k$ , then by equation (1.9) we have

$$v_1 \wedge v_2 \cdot d_H \varphi = 2[v_1, v_2] \cdot \varphi - d_H(v_1 \wedge v_2 \cdot \varphi) - 2v_1 \cdot d_H(v_2 \cdot \varphi) + 2v_2 \cdot d_H(v_1 \cdot \varphi)$$

but the right hand side is, by inductive hypothesis, in  $\oplus_{j \geq k+1} \mathcal{U}^j$ . Therefore, so is  $v_2 \wedge v_1 \cdot d_H \varphi$ . Since  $v_1$  and  $v_2$  are sections of  $L$ , we conclude that  $d_H \varphi \in \oplus_{j \geq k-1} \mathcal{U}^j$ .

In order to finish the proof of the converse, we remark that conjugation swaps  $\mathcal{U}^{\pm k}$ , but preserves  $d_H$ , as it is a real operator, i.e.,  $d_H\varphi = \overline{d_H\bar{\varphi}}$ . Thus, for  $\varphi \in \mathcal{U}^k$ ,  $\bar{\varphi} \in \mathcal{U}^{-k}$  and, using (1.13),

$$d_H\varphi \in \bigoplus_{j \geq k-1} \mathcal{U}^j \quad d\varphi = \overline{d_H\bar{\varphi}} \in \bigoplus_{j \leq k+1} \mathcal{U}^j.$$

Showing that  $d_H\varphi \in \mathcal{U}^{k-1} \oplus \mathcal{U}^k \oplus \mathcal{U}^{k+1}$ . Finally, if  $\varphi \in \mathcal{U}^k$  is an even/odd form, then all the elements in  $\mathcal{U}^k$  are even/odd whereas the elements of  $\mathcal{U}^{k-1} \oplus \mathcal{U}^{k+1}$  are odd/even. As  $d_H$  swaps the parity of the degree of forms,  $d_H\varphi$  is odd/even and hence has no  $\mathcal{U}^k$  component.  $\square$

So, on a generalized complex manifold  $M$  we can define the operators

$$\begin{aligned} \partial : \mathcal{U}^k &\longrightarrow \mathcal{U}^{k+1}; \\ \bar{\partial} : \mathcal{U}^k &\longrightarrow \mathcal{U}^{k-1}; \end{aligned}$$

as the projections of  $d_H$  onto each of these factors. We also define  $d^{\mathcal{J}} = -i(\partial - \bar{\partial})$ .

Similarly to the operator  $d^c$  from complex geometry, we can find other expressions for  $d^{\mathcal{J}}$  based on the action of the generalized complex structure on forms. One can easily check that if we consider the Lie group action of  $\mathcal{J}$ , i.e,  $\mathcal{J}$  acts on  $\mathcal{U}^k$  as multiplication by  $i^k$ , then

$$d^{\mathcal{J}} = \mathcal{J}^{-1}d_H\mathcal{J}.$$

And if one considers the Lie algebra action, then

$$d^{\mathcal{J}} = [d_H, \mathcal{J}].$$

As a consequence of Example 1.21 it is clear that in the complex case,  $\partial$  and  $\bar{\partial}$  are just the standard operators denoted by the same symbols and  $d^{\mathcal{J}} = d^c$ . In the symplectic case,  $d^{\mathcal{J}}$  corresponds to Koszul's *canonical homology derivative* [46] introduced in the context of Poisson manifolds and studied by Brylinski [9], Mathieu [53], Yan [68], Merkulov [55] and others [26, 41] in the symplectic setting.

In the symplectic case, the operators  $\partial$  and  $\bar{\partial}$  are related to  $d$  and  $d^{\mathcal{J}}$  also in a more subtle way. Recall from Example 1.10 that for a symplectic structure we have a map  $\Phi : \wedge^{\bullet}T^*M \longrightarrow \wedge^{\bullet}T^*M$  such

that  $\Phi(\wedge^k T^*M) = U^{n-k}$ . The operators  $\partial$ ,  $\bar{\partial}$ ,  $d$  and  $d^{\mathcal{J}}$  and the map  $\Phi$  are related by (see [16])

$$\bar{\partial}\Phi(\alpha) = \Phi(d\alpha) \quad 2i\partial\Phi(\alpha) = \Phi(d^{\mathcal{J}}\alpha).$$

**Example 1.26.** If a generalized complex structure induces a splitting of  $\wedge^{\bullet}T^*M$  into subspaces  $U^k$ , then, according to Example 1.11, a  $B$ -field transform of this structure will induce a decomposition into  $e^B U^k$ . As  $B$  is a closed form, for  $v \in U^k$  we have:

$$d(e^B v) = e^B dv = e^B \partial v + e^B \bar{\partial} v \in e^B \mathcal{U}^{k+1} + e^B \mathcal{U}^{k-1},$$

hence the corresponding operators for the  $B$ -field transform,  $\partial_B$  and  $\bar{\partial}_B$ , are given by

$$\partial_B = e^B \partial e^{-B}; \quad \bar{\partial}_B = e^B \bar{\partial} e^{-B}.$$

### 1.3.1 The differential graded algebra $(\Omega^{\bullet}(\bar{L}), d_L)$

A peculiar characteristic of the operators  $\partial$  and  $\bar{\partial}$  introduced last section is that they are not derivations, i.e., they do not satisfy the Leibniz rule. There is, however, another differential complex associated to a generalized complex structure for which the differential is a derivation. We explain this in this section.

As we mentioned before, the Courant bracket does not satisfy the Jacoby identity, and instead we have

$$\text{Jac}(e_1, e_2, e_3) = \frac{1}{3}d(\langle [e_1, e_2], e_3 \rangle + c.p.).$$

However, the identity above also shows that the Courant bracket induces a Lie bracket when restricted to sections of any involutive isotropic space  $L$ . This Lie bracket together with the projection  $\pi_T : L \rightarrow TM$ , makes  $L$  into a Lie algebroid and allows us to define a differential  $d_L$  on  $\Omega^{\bullet}(L^*) = \Gamma(\wedge^{\bullet}L^*)$  making it into a differential graded algebra (DGA). This is analogous to the way the Lie bracket of vector fields determines the exterior derivative  $d$ . If  $L$  is the  $i$ -eigenspace of a generalized complex structure, then the natural pairing gives an isomorphism  $L^* \cong \bar{L}$  and with this identification  $(\Omega^{\bullet}(\bar{L}), d_L)$  is a DGA.



If a generalized complex structure has type zero over  $M$ , i.e., is of symplectic type, then  $\pi : L \xrightarrow{\cong} T_{\mathbb{C}}M$  is an isomorphism and the Courant bracket on  $\Gamma(L)$  is mapped to the Lie bracket on  $\Gamma(T_{\mathbb{C}}M)$ . Therefore, in this particular case,  $(\Omega^{\bullet}(\bar{L}), d_L)$  and  $(\Omega_{\mathbb{C}}^{\bullet}(M), d)$  are isomorphic DGAs.

Further, recall that the choice of a nonvanishing local section  $\rho$  of the canonical bundle gives an isomorphism of Clifford modules:

$$\phi : \Omega^{\bullet}(\bar{L}) \longrightarrow \Omega_{\mathbb{C}}^{\bullet}(M); \quad \phi(s \cdot \sigma) = s \cdot \rho.$$

With these choices, the operators  $\bar{\partial}$  and  $d_L$  are related by

$$\bar{\partial}\phi(\alpha) = \phi(d_L\alpha) + (-1)^{|\alpha|}\alpha \cdot d_H\rho.$$

In the particular case when there is a nonvanishing global holomorphic section  $\rho$  we can define  $\phi$  globally and have

$$\bar{\partial}\phi(\alpha) = \phi(d_L\alpha).$$

## 1.4 Deformations of generalized complex structures

In this section we state part of Gualtieri's deformation theorem for generalized complex structures. The first observation is that small deformations of a generalized complex structure correspond to small deformations of its  $+i$ -eigenspace,  $L$ , and those can be described as the graph of a linear map  $\varepsilon \in \text{Hom}(L, \bar{L}) \cong L^* \otimes \bar{L} \cong \otimes^2 \bar{L}$ .

From the linear algebra point of view, the requirement that  $L^{\varepsilon}$  is isotropic amounts to asking for  $\varepsilon \in \wedge^2 \bar{L} \subset \otimes^2 \bar{L}$ . The condition  $L \cap \bar{L} = \{0\}$ , is open, so small deformations won't spoil it. Observe that the deformed structure has  $i$ -eigenspace  $L^{\varepsilon} = e^{\varepsilon} \cdot L$  and  $-i$ -eigenspace  $\bar{L}^{\varepsilon} = e^{\bar{\varepsilon}} \cdot \bar{L}$ .

So, the only nontrivial part of when deciding whether  $L^{\varepsilon}$  is a generalized complex structure is the integrability condition. Drawing on a result of Liu, Weinstein and Xu [51], Gualtieri established the following deformation theorem.




Figure 1.1: Small deformations of a generalized complex structure are given by the graph of an element  $\varepsilon \in \wedge^2 \bar{L}$

**Theorem 1.27** (Gualtieri [34], Theorem 5.4). *An element  $\varepsilon \in \wedge^2 \bar{L}$  gives rise to a deformation of generalized complex structures if and only if  $\varepsilon$  is small enough and satisfies the Maurer–Cartan equation*

$$d_L \varepsilon + \frac{1}{2} [[\varepsilon, \varepsilon]] = 0.$$

The deformed generalized complex structure is given by

$$L^\varepsilon = (\text{Id} + \varepsilon)L \quad \bar{L}^\varepsilon = (\text{Id} + \bar{\varepsilon})\bar{L}.$$

In the complex case, a bivector  $\varepsilon \in \wedge^{2,0} TM \subset \wedge^2 \bar{L}$  gives rise to a deformation only if each of the summands vanish, i.e.,  $\bar{\partial}\varepsilon = 0$  ( $\varepsilon$  is holomorphic) and  $[[\varepsilon, \varepsilon]] = 0$  ( $\varepsilon$  is Poisson).

*Remark:* The theorem above is only concerned with which 2-forms  $\varepsilon \in \Gamma(\wedge^2 \bar{L})$  give rise to integrable generalized complex structures, but it does not tackle the problem of determining which ones give

rise to *trivial deformations*, i. e., deformations which are obtained by  $B$ -field transforms and diffeomorphisms. We shall not discuss this here and instead refer to Gualtieri's thesis.

**Example 1.28.** Consider  $\mathbb{C}^2$  with its standard complex structure and let  $\varepsilon = z_1 \partial_{z_1} \wedge \partial_{z_2}$ . One can easily check that  $\varepsilon$  is a holomorphic Poisson bivector, hence we can use  $\varepsilon$  to deform the complex structure on  $\mathbb{C}^2$ . According to Example 1.5, the canonical bundle of the deformed structure is given by

$$e^\varepsilon \cdot dz_1 \wedge dz_2 = z_1 + dz_1 \wedge dz_2.$$

One can readily recognize this as the generalized complex structure from Example 1.24. This illustrates the fact that in 2 complex dimensions the zeros of the holomorphic bivector correspond to the type-change points in the deformed structure.

**Example 1.29.** Any holomorphic bivector  $\varepsilon$  on a complex surface  $M$  is also Poisson, as  $[\varepsilon, \varepsilon] \in \wedge^{3,0}TM = \{0\}$ , and hence gives rise to a deformation of generalized complex structures. The deformed generalized complex structure will be symplectic outside the divisor representing  $c_1(M)$  where the bivector vanishes. At the points where  $\varepsilon = 0$  the deformed structure agrees with the original complex structure.

**Example 1.30.** Let  $M^{4n}$  be a hyperkähler manifold with Kähler forms  $\omega_I, \omega_J$  and  $\omega_K$ . According to the Kähler structure  $(\omega_I, I)$ ,  $(\omega_J + i\omega_K)$  is a closed holomorphic 2-form and  $(\omega_J + i\omega_K)^n$  is a holomorphic volume form. Therefore these generate a holomorphic Poisson bivector  $\Lambda \in \wedge^{2,0}TM$  by

$$\Lambda \cdot (\omega_J + i\omega_K)^n = n(\omega_J + i\omega_K)^{n-1}.$$

The deformation of the complex structure  $I$  by  $t\Lambda$  is given by

$$e^{t\Lambda}(\omega_J + i\omega_K)^n = t^n e^{\frac{\omega_J + i\omega_K}{t}}.$$

which interpolates between the complex structure  $I$  and the  $B$ -field transform of the symplectic structure  $\omega_K$  as  $t$  varies from 0 to 1.

## 1.5 Submanifolds

In this section we introduce two special types of submanifolds of a generalized complex manifold. The first type are the *generalized Lagrangians* introduced by Gualtieri [34]. This class of submanifolds comprises complex submanifolds from complex geometry and Lagrangian submanifolds from symplectic geometry. Generalized Lagrangians are intimately related to branes [34, 42, 70]. The second type of submanifolds consists of those which inherit a generalized complex structure from the original manifold. These correspond to one of the definitions of submanifolds introduced by Ben-Bassat and Boyarchenko [6] and are the analogue of symplectic submanifolds from symplectic geometry.

In order to understand generalized complex submanifolds it is desirable to understand how to restrict Courant algebroids to submanifolds. Given a Courant algebroid  $\mathcal{E}$  over a manifold  $M$  and a submanifold  $\iota : N \hookrightarrow M$  there are two natural bundles one can form over  $N$ . The first is  $\iota^*\mathcal{E}$ , the pull back of  $\mathcal{E}$  to  $N$ . The pairing on  $\mathcal{E}$  induces a pairing on  $\iota^*\mathcal{E}$ , however the same is not true about the Courant bracket: even if a section vanishes over  $N$ , it may bracket nonzero with a nonvanishing section.

**Exercise 1.31.** Show that if  $e_1, e_2 \in \Gamma(\mathcal{E})$ ,  $\pi(\iota^*e_1) \in \Gamma(TN)$  and  $\iota^*e_2 = 0$ , then  $\iota^*[[e_1, e_2]]$  is not necessarily zero but lies in  $\mathcal{N}^* = \text{Ann}(TN) \subset T^*M$ , the conormal bundle of  $N$ .

The second bundle, called the *restricted Courant algebroid*, and denoted by  $\mathcal{E}|_N$ , is a Courant algebroid, as the name suggests. It is defined by

$$\mathcal{E}|_N = \frac{\mathcal{N}^{*\perp}}{\mathcal{N}^*} = \frac{\{e \in \iota^*\mathcal{E} : \pi(e) \in TN\}}{\text{Ann}(TN)}.$$

The bracket is defined using the Courant bracket on  $\mathcal{E}$ : according to Exercise 1.31, the ambiguity of the bracket on  $\mathcal{N}^{*\perp}$  lies in  $\mathcal{N}^*$ , hence the bracket on  $\mathcal{E}|_N$  is well defined. Since  $\mathcal{N}^* \subset \mathcal{N}^{*\perp}$  is the null space of the pairing restricted to  $\mathcal{N}^{*\perp}$ , we obtain a well defined nondegenerate pairing on  $\mathcal{E}|_N$ . These bracket and pairing make  $\mathcal{E}|_N$  a Courant algebroid. If  $\mathcal{E}$  is split and has curvature  $H$ ,  $\mathcal{E}|_N$  is naturally isomorphic to  $TN \oplus T^*N$  endowed with the  $\iota^*H$ -Courant bracket.

Using these two bundles we can define two types of generalized complex submanifolds.

**Definition 1.32.** Given a generalized complex structure  $\mathcal{J}$  on a Courant algebroid  $\mathcal{E}$  over a manifold  $M$ , a *generalized Lagrangian* is a submanifold  $\iota : N \rightarrow M$  together with a maximal isotropic subbundle  $\tau_N \subset \iota^*\mathcal{E}$  invariant under  $\mathcal{J}$  such that  $\pi(\tau_N) = TN$  and if  $\iota^*e_1, \iota^*e_2 \in \Gamma(\tau_N)$  then  $\iota^*[[e_1, e_2]] \in \Gamma(\tau_N)$ .

Since  $\tau_N$  is maximal isotropic and  $\pi(\tau_N) = TN$ , it follows that  $\tau_N$  sits in an exact sequence

$$0 \rightarrow \mathcal{N}^* \rightarrow \tau_N \rightarrow TN \rightarrow 0.$$

Since  $\mathcal{N}^* \subset \tau_N$ , it makes sense to ask for  $\tau_N$  to be closed under the bracket on  $\iota^*\mathcal{E}$ , even though this bracket is not well defined, since the indeterminacy lies in  $\mathcal{N}^*$ .

**Exercise 1.33.** Show that if  $\mathcal{E}$  is split, then there is  $F \in \Omega^2(N)$  such that

$$\tau_N = (Id + F) \cdot TN \oplus \mathcal{N}^* = \{X + \xi \in TN \oplus T^*M : \xi|_{TN} = i_X F\}. \quad (1.14)$$

Show that  $\tau_N$  is closed under the bracket if and only if  $dF = \iota^*H$ , where  $H$  is the curvature of the splitting. Therefore a necessary condition for a manifold to be a submanifold is that  $\iota^*[H] = 0$ .

Using this exercise we obtain an alternative description of a generalized Lagrangian for a split Courant algebroid: it is a submanifold  $N$  with a 2-form  $F \in \Omega^2(N)$  such that  $dF = \iota^*H$  and  $\tau_N$ , as defined in (1.14), is invariant under  $\mathcal{J}$ .

For the second definition of submanifold we observe that there is a natural way to transport a Dirac structure  $D$  on  $\mathcal{E}$  to a Dirac structure on  $\mathcal{E}|_N$ :

$$D_{red} = \frac{D \cap \mathcal{N}^{*\perp} + \mathcal{N}^*}{\mathcal{N}^*}.$$

The distribution  $D_{red}$  is a Dirac structure whenever it is smooth and it is called the *pull-back* of  $D$ . So, if  $L$  is the  $i$ -eigenspace of a generalized complex structure  $\mathcal{J}$  on  $\mathcal{E}$ ,  $L_{red}$ , the pull back of  $L$ , is a Dirac structure on  $\mathcal{E}|_N$ .

**Definition 1.34.** A *generalized complex submanifold* is a submanifold  $\iota : N \rightarrow M$  for which  $\underline{L_{red}}$  determines a generalized complex structure on  $\mathcal{E}|_N$ , i.e.,  $L_{red} \cap \overline{L_{red}} = \{0\}$ .

One can check that  $L_{red}$  is a generalized complex structure if and only if it is smooth and satisfies (c.f. Lemma 3.19)

$$\mathcal{J}\mathcal{N}^* \cap \mathcal{N}^{*\perp} \subset \mathcal{N}^*.$$

Some particular cases when the above holds are when  $\mathcal{N}^*$  is  $\mathcal{J}$  invariant, i.e.,  $\mathcal{J}\mathcal{N}^* = \mathcal{N}^*$  or when the natural pairing is nondegenerate on  $\mathcal{J}\mathcal{N}^* \times \mathcal{N}^*$ .

**Example 1.35** (Gualtieri [34], Example 7.8). In this example we describe generalized Lagrangians of a symplectic manifold. So, the starting point is the split Courant algebroid  $TM \oplus T^*M$  with  $H = 0$  and generalized complex structure given by Example 1.3. Since the Courant algebroid is split, we can use the description of  $\tau_N$  in terms of  $TN$  and a 2-form  $F$ . In this case, as was observed by Gualtieri, the definition of generalized Lagrangian agrees with the A-branes of Kapustin and Orlov [42].

We claim that if  $(N, F)$  is a generalized Lagrangian, then  $N$  is a coisotropic submanifold. Also, both  $F$  and, obviously,  $\omega|_N$  are annihilated by the distribution  $TN^\omega = \omega^{-1}(\mathcal{N}^*)$ . In the quotient  $V = TN/TN^\omega$ , there is a complex structure induced by  $\omega^{-1}F$  and  $(F + i\omega)$  is a  $(2, 0)$ -form whose top power is a volume element in  $\wedge^{k, 0}V$ . Finally, if  $F = 0$ , then  $N$  is just a Lagrangian submanifold of  $M$ .

To prove that  $N$  is coisotropic, we have to prove that  $\omega^{-1}(\mathcal{N}^*) \subset TN$ . This is a simple consequence of the fact that  $\mathcal{N}^* \subset \tau_N$  and  $\tau_N$  is  $\mathcal{J}$  invariant, hence

$$\mathcal{J}\mathcal{N}^* = -\omega^{-1}\mathcal{N}^* \in \tau_N,$$

showing that  $N$  is coisotropic.

To show that  $F$  is annihilated by  $TN^\omega$ , we choose a local extension,  $B$ , of  $F$  to  $\Omega^2(M)$ , so that for  $X \in TN$ ,  $X + B(X) \in \tau_N$ . Since  $\tau_N$  is  $\mathcal{J}$  invariant,

$$\mathcal{J}(X + B(X)) = -\omega^{-1}B(X) + \omega(X) \in \tau_N,$$

and hence  $\omega^{-1}B(X) \in TN$ , which implies that  $B(X)$  vanishes on  $TN^\omega = \omega^{-1}(\mathcal{N}^*)$  since

$$0 = \langle \omega^{-1}B(X), \mathcal{N}^* \rangle = \langle B(X), -\omega^{-1}(\mathcal{N}^*) \rangle.$$

Hence  $F$  is also annihilated by the distribution  $TN^\omega$ .

To find the complex structure on the quotient  $TN/TN^\omega$  we take  $X \in TN$  and apply  $\mathcal{J}$  to  $X + B(X) \in \tau_N$ . Invariance implies that

$$-\omega^{-1}B(X) + \omega(X) \in \tau_N,$$

which is the same as  $-F \circ \omega^{-1} \circ F(X)|_N = \omega(X)|_N$ . In  $TN/TN^\omega$ , there is an inverse  $\omega^{-1}$  and hence, in the quotient, we have the identity

$$-X = (\omega^{-1}F)^2(X),$$

showing that  $\omega^{-1}F$  induces a complex structure on  $TN/TN^\omega$ .

For an  $X \in \wedge^{0,1}(TN/TN^\omega)$  we have  $\omega^{-1}F(X) = -iX$ . Applying  $\omega$  we get  $F(X, \cdot) + i\omega(X, \cdot) = 0$  and hence  $F + i\omega$  is annihilated by  $\wedge^{0,1}(TN/TN^\omega)$  and thus is a  $(2, 0)$ -form. Finally, for  $X = X_1 + iX_2 \in \wedge^{1,0}(TN/TN^\omega)$ , as before we obtain  $(F - i\omega)(X, \cdot) = 0$ , which spells out as

$$F(X_1, \cdot) = -\omega(X_2, \cdot) \quad \text{and} \quad F(X_2, \cdot) = \omega(X_1, \cdot),$$

and therefore  $(F + i\omega)(X, \cdot) = -2\omega(X_2, \cdot) + 2\omega(X_1, \cdot) \neq 0$ , as  $\omega$  is nondegenerate in  $TN/TN^\omega$ . Thus  $F + i\omega$  is a nondegenerate  $2, 0$ -form.

If  $F$  vanishes,  $0$  is a complex structure on  $TN/TN^\omega$  which must therefore be the trivial vector space and hence  $N$  is Lagrangian.

**Exercise 1.36.** Show that a generalized Lagrangian of a complex manifold is a complex submanifold with a closed form  $F \in \Omega^{1,1}(M)$ .

**Exercise 1.37.** Show that generalized complex submanifolds of a complex manifold are complex while those of a symplectic manifold are symplectic submanifolds.

## 1.6 Examples

In this section we give some interesting examples of generalized complex structures. In the first example we find generalized complex

structures on symplectic fibrations and, in the second, on Lie algebras. The last example consists of a surgery procedure for symplectic manifolds which produces generalized complex structures on Courant algebroids over a topologically distinct manifold and whose characteristic class is not necessarily trivial.

### 1.6.1 Symplectic fibrations

The differential form description of a generalized complex structure furnishes also a very pictorial one around *regular points*. Indeed, in this case one can choose locally a closed form  $\rho$  defining the structure. Then the integrability condition tells us that  $\Omega \wedge \overline{\Omega}$  is a real closed form and therefore the distribution  $\text{Ann}(\Omega \wedge \overline{\Omega}) \subset TM$  is an integrable distribution. The algebraic condition  $\Omega \wedge \overline{\Omega} \wedge \omega^{n-k} \neq 0$  implies that  $\omega$  is nondegenerate on the leaves and the integrability condition  $\Omega \wedge d\omega = 0$ , that  $\omega$  is closed when restricted to these leaves. Therefore, around a regular point, the generalized complex structure furnishes a natural symplectic foliation, and further, the space of leaves has a natural complex structure given by  $\Omega$ .

This suggests that symplectic fibrations should be a way to construct nontrivial examples of generalized complex structures. Next we see that Thurston's argument for symplectic fibrations [63, 54] can also be used in the generalized complex setting.

**Theorem 1.38.** *Let  $M$  be a generalized complex manifold with the  $H$ -Courant bracket. Let  $\pi : P \rightarrow M$  be a symplectic fibration with compact fiber  $(F, \omega)$ . Assume that there is  $a \in H^2(P)$  which restricts to the cohomology class of  $\omega$  on each fiber. Then  $P$  admits a generalized complex structure with the  $\pi^*H$ -Courant bracket.*

*Proof.* The usual argument using partitions of unit shows that we can find a closed 2-form  $\tau$  representing the cohomology class  $a$  such that  $\tau|_F = \omega$  on each fiber. If  $K$  is the canonical bundle of the generalized complex structure on  $M$ , then we claim that, for  $\varepsilon$  small enough, the subbundle  $K_P = e^{i\varepsilon\tau} \wedge \pi^*K \subset \Omega_{\mathbb{C}}^{\bullet}(P)$  determines a generalized complex structure on  $P$  with the  $\pi^*H$ -Courant bracket.

Let  $U_{\alpha}$  be a covering of  $B$  where we have trivializations  $\rho_{\alpha}$  of  $K$ . Then the forms  $e^{i\varepsilon\tau} \wedge \pi^*\rho_{\alpha}$  are nonvanishing local sections of



$K_P$ . Since  $\tau$  is nondegenerate on the vertical subspaces  $\text{Ker } \pi_*$  it determines a field of horizontal subspaces

$$\text{Hor}_x = \{X \in T_x P : \tau(X, Y) = 0, \forall Y \in T_x F\}.$$

The subspace  $\text{Hor}_x$  is a complement to  $T_x F$  and isomorphic to  $T_{\pi(x)} M$  via  $\pi_*$ . Also, denoting by  $(\cdot, \cdot)_M$  the Mukai pairing on  $M$ ,  $(\rho_\alpha, \overline{\rho_\alpha})_M \neq 0$  and hence pulls back to a volume form on  $\text{Hor}$ . Therefore, for  $\varepsilon$  small enough,

$$(e^{i\varepsilon\tau} \wedge \pi^* \rho_\alpha, e^{-i\varepsilon\tau} \wedge \pi^* \overline{\rho_\alpha})_P = (\varepsilon\tau)^{\dim(F)} \wedge \pi^*(\rho_\alpha, \overline{\rho_\alpha})_M \neq 0,$$

and  $e^{i\varepsilon\tau} \wedge \pi^* \rho_\alpha$  is of the right algebraic type. Finally, from the integrability condition for  $\rho_\alpha$ , there are  $X_\alpha, \xi_\alpha$  such that

$$d_H \rho_\alpha = (X_\alpha + \xi_\alpha) \rho_\alpha.$$

If we let  $X_\alpha^{\text{hor}} \in \text{Hor}$  be the horizontal vector projecting down to  $X_\alpha$ , then the following holds on  $\pi^{-1}(U_\alpha)$ :

$$d_{\pi^* H}(e^{i\varepsilon\tau} \wedge \pi^* \rho_\alpha) = (X_\alpha^{\text{hor}} + \pi^* \xi_\alpha - X_\alpha^{\text{hor}} \lrcorner i\varepsilon\tau) e^{i\varepsilon\tau} \wedge \pi^* \rho_\alpha,$$

showing that the induced generalized complex structure is integrable.  $\square$

Several cases for when the conditions of the theorem are fulfilled have been studied for symplectic manifolds and many times purely topological conditions on the base and on the fiber are enough to ensure that the hypotheses hold. We give the following two examples adapted from McDuff and Salamon [54], Chapter 6.

**Theorem 1.39.** *Let  $\pi : P \rightarrow M$  be a symplectic fibration over a compact generalized complex base with fiber  $(F, \omega)$ . If the first Chern class of  $TF$  is a nonzero multiple of  $[\omega]$ , then the conditions of Theorem 1.38 hold. In particular, any oriented surface bundle can be given a symplectic fibration structure and, if the fibers are not tori, the total space has a generalized complex structure.*

**Theorem 1.40.** *A symplectic fibration with compact and 1-connected fiber and a compact twisted generalized complex base admits a twisted generalized complex structure.*

### 1.6.2 Nilpotent Lie algebras

Another source of examples comes from considering left invariant generalized complex structures on Lie groups, which is equivalent to consider integrable linear generalized complex structures on their Lie algebra  $\mathfrak{g}$ . In this case, one of the terms in the  $H$ -Courant bracket always vanishes and we have

$$[[X + \xi, Y + \eta]] = [X, Y] + \mathcal{L}_X \eta - \mathcal{L}_Y \xi + i_Y i_X H,$$

which is a Lie bracket on  $\mathfrak{g} \oplus \mathfrak{g}^*$ .

Hence the search for a generalized complex structure on a Lie algebra  $\mathfrak{g}$  amounts to finding a complex structure on  $\mathfrak{g} \oplus \mathfrak{g}^*$  orthogonal with respect to the natural pairing. Therefore, given a fixed Lie algebra, finding a generalized complex structure on it or proving it does not admit any such structure is only a matter of perseverance.

In joint work with Gualtieri, the author carried out this task studying generalized complex structures on nilpotent Lie algebras. There we proved

**Theorem 1.41** (Cavalcanti–Gualtieri [19]). *Every 6-dimensional nilpotent Lie algebra has a generalized complex structure.*

A classification of which of those algebras had complex or symplectic structures had been carried out before by Salamon [57] and Goze and Khakimdjanov [32] and 5 nilpotent Lie algebras don't have either. Therefore these results gave the first nontrivial instances of generalized complex structures on spaces which admitted neither (left invariant) complex or symplectic structures.

Similar work was also done for 4-dimensional solvable Lie algebras in [24]. In contrast, there they show that the 4-dimensional solvable Lie algebras admitting generalized complex structures coincide with those admitting either complex or symplectic structures.

### 1.6.3 A surgery

One of the features of generalized complex structures is that they don't necessarily have constant type along the manifold, as we saw in Example 1.24. In four dimensions, this is one of the main features

distinguishing generalized complex structures from complex or symplectic structures and was used to produce some interesting examples in [20] by means of a surgery.

The idea of the surgery is to replace a neighborhood  $U$  of a symplectic 2-torus  $T$  with trivial normal bundle on a symplectic manifold  $(M, \sigma)$  by  $D^2 \times T^2$  with the generalized complex structure from Example 1.24 using a symplectomorphism which is a nontrivial diffeomorphism of  $\partial U \cong T^3$ . This surgery is a particular case a  $C^\infty$  logarithmic transformation, a surgery studied by Gompf and Mrowka in [31].

**Theorem 1.42** (Cavalcanti–Gualtieri [20]). *Let  $(M, \sigma)$  be a symplectic 4-manifold,  $T \hookrightarrow M$  be a symplectic 2-torus with trivial normal bundle and tubular neighbourhood  $U$ . Let  $\psi : S^1 \times T^2 \rightarrow \partial U \cong S^1 \times T^2$  be the map given on standard coordinates by*

$$\psi(\theta_1, \theta_2, \theta_3) = (\theta_3, \theta_2, -\theta_1).$$

Then

$$\tilde{M} = M \setminus U \cup_\psi D^2 \times T^2,$$

admits a generalized complex structure with type change along a 2-torus, and which is integrable with respect to a 3-form  $H$ , such that  $[H]$  is the Poincaré dual to the circle in  $S^1 \times T^2$  preserved by  $\psi$ . If  $M$  is simply connected and  $[T] \in H^2(M, \mathbb{Z})$  is  $k$  times a primitive class, then  $\pi_1(\tilde{M}) = \mathbb{Z}_k$ .

*Proof.* By Moser's theorem, symplectic structures with the same volume on an oriented compact surface are isomorphic. Hence, after rescaling, we can assume that  $T$  is endowed with its standard symplectic structure. Therefore, by Weinstein's neighbourhood theorem [65], the neighborhood  $U$  is symplectomorphic to  $D^2 \times T^2$  with standard symplectic form:

$$\sigma = \frac{1}{2} d\tilde{r}^2 \wedge d\tilde{\theta}_1 + d\tilde{\theta}_2 \wedge d\tilde{\theta}_3.$$

Now consider the symplectic form  $\omega$  on  $D^2 \setminus \{0\} \times T^2$  from Example 1.24:

$$\omega = d \log r \wedge d\theta_3 + d\theta_1 \wedge d\theta_2.$$

The map  $\psi : (D^2 \setminus D_{1/\sqrt{\epsilon}}^2 \times T^2, \omega \longrightarrow (D^2 \setminus \{0\} \times T^2, \sigma)$  given by

$$\psi(r, \theta_1, \theta_2, \theta_3) = (\sqrt{\log er^2}, \theta_3, \theta_2, -\theta_1)$$

is a symplectomorphism.

Let  $B$  be the 2-form defined by (1.12) on  $D^2 \setminus D_{1/\sqrt{\epsilon}}^2 \times T^2$ , and choose an extension  $\tilde{B}$  of  $\psi^{-1*}B$  to  $M \setminus T$ . Therefore  $(M \setminus T, \tilde{B} + i\sigma)$  is a generalized complex manifold of type 0, integrable with respect to the  $d\tilde{B}$ -Courant bracket.

Now the surgery  $\tilde{M} = M \setminus T \cup_{\psi} D^2 \times T^2$  obtains a generalized complex structure since the gluing map  $\psi$  satisfies  $\psi^*(\tilde{B} + i\sigma) = B + i\omega$ , therefore identifying the generalized complex structures on  $M \setminus T$  and  $D^2 \times T^2$  over the annulus where they are glued together. Therefore, the resulting generalized complex structure exhibits type change along the 2-torus coming from the central fibre of  $D^2 \times T^2$ . This structure is integrable with respect to  $H = d\tilde{B}$ , which is a globally defined closed 3-form on  $\tilde{M}$ .

The 2-form  $\tilde{B}$  can be chosen so that it vanishes outside a larger tubular neighbourhood  $U'$  of  $T$ , so that  $H = d\tilde{B}$  has support in  $U' \setminus U$  and has the form

$$H = f'(r)dr \wedge d\theta_1 \wedge d\theta_3,$$

for a smooth bump function  $f$  such that  $f|_U = 1$  and vanishes outside  $U'$ . Therefore, we see that  $H$  represents the Poincaré dual of the circle parametrized by  $\theta_2$ , as required.

The last claim is a consequence of van Kampen's theorem and that  $H^2(M, \mathbb{Z})$  is spherical, as  $M$  is simply connected.  $\square$

**Example 1.43.** Given an elliptic K3 surface, one can perform the surgery above along one of the  $T^2$  fibers. In [31], Gompf and Mrowka show that the resulting manifold is diffeomorphic to  $3\mathbb{C}P^2 \# 19\overline{\mathbb{C}P^2}$ . Due to Taubes's results on Seiberg–Witten invariants [62] and Kodaira's classification of complex surfaces [45], we know this manifold does not admit symplectic or complex structures therefore providing the first example of generalized complex manifold without complex or symplectic structures.

## Chapter 2

# Generalized metric structures

In this chapter we present metrics on Courant algebroids as introduced in [34, 67] and further developed in [35]. Since a Courant algebroid is endowed with a natural pairing, one has to place a compatibility condition between metric and pairing. This is done by defining that a *generalized metric* is a self adjoint, orthogonal endomorphism  $\mathcal{G} : \mathcal{E} \longrightarrow \mathcal{E}$  such that, for  $v \in \mathcal{E} \setminus \{0\}$ ,

$$\langle \mathcal{G}v, v \rangle > 0.$$

A generalized metric on a split Courant algebroid gives rise to a Hodge star-like operator on forms. Further, for a given generalized metric, there is a natural splitting of any exact Courant algebroid as the sum of  $T^*M$  with its metric orthogonal complement,  $(T^*M)^{\mathcal{G}}$ . If this splitting is chosen, the star operator coincides with the usual Hodge star, while in general it differs from it by nonclosed  $B$ -field transforms.

Similarly to the complex case, one can ask for a generalized metric to be compatible with a given generalized complex structure. There always are such metrics. Whenever a metric compatible with a generalized complex structure is chosen, we automatically get a second generalized complex structure which is not integrable in general. By

studying Hodge theory on generalized complex manifolds we obtain Serre duality for the operator  $\bar{\partial}$ . When both of the generalized complex structures are integrable, we obtain a generalized Kähler structure.

The compatibility between a metric and a generalized complex structure is used to the full in the case of a generalized Kähler structure, for Gualtieri proved that in a generalized Kähler manifold a number of Laplacians coincide [35] furnishing Hodge identities for those manifolds. These are powerful results which have implications for a generalized Kähler manifold similar to the formality theorem for Kähler manifolds [25], as made explicit in [15]. Moreover these identities are a key fact for the proof of smoothness of the moduli space of generalized Kähler structures [47].

After introducing the generalized metric, this chapter follows closely [35] and gives some applications of the results therein. In the first section we introduce the concept of generalized metric on a vector space and investigate the consequences of the compatibility of this metric with a linear generalized complex structure. In Section 2.2 we do things over a manifold with the requirement that the generalized complex structure involved is integrable. We also state Gualtieri's theorem relating generalized Kähler structures to bihermitian structures. In Section 2.3 we study Hodge theory on a generalized Kähler manifold and in Section 2.4 we give an application of those results.

## 2.1 Linear algebra of the metric

### 2.1.1 Generalized metric

The concept of generalized metric on  $\mathcal{D}V$ , the double of a vector space  $V$ , was introduced by Gualtieri [34] and Witt [67] and was further studied by Gualtieri in connection with generalized complex structures in [35]. Following Gualtieri's exposition, a *generalized metric* is an orthogonal, self adjoint operator  $\mathcal{G} : \mathcal{D}V \rightarrow \mathcal{D}V$  such that

$$\langle \mathcal{G}e, e \rangle > 0 \text{ for } e \in \mathcal{D}V \setminus \{0\}.$$

Using symmetry and orthogonality we see that

$$\mathcal{G}^2 = \mathcal{G}\mathcal{G}^t = \mathcal{G}\mathcal{G}^{-1} = \text{Id}.$$

Hence  $\mathcal{G}$  splits  $\mathcal{D}V$  into its  $\pm 1$ -eigenspaces,  $C_{\pm}$ , which are orthogonal subspaces of  $\mathcal{D}V$  where the pairing is  $\pm$ -definite. Therefore  $C_{\pm}$  are maximal and, since  $V^* \subset \mathcal{D}V$  is isotropic, the projections  $\pi : C_{\pm} \rightarrow V$  are isomorphisms. Conversely, prescribing orthogonal spaces  $C_{\pm}$  where the pairing is  $\pm$ -definite defines a metric  $\mathcal{G}$  by letting  $\mathcal{G} = \pm 1$  on  $C_{\pm}$ .

A generalized metric induces a metric on the underlying vector space. This is obtained using the isomorphism  $\pi : C_+ \rightarrow V$  and defining

$$g(X, Y) = \langle \pi^{-1}(X), \pi^{-1}(Y) \rangle.$$

One can alternatively define a metric using  $C_-$ , but this renders the same metric on  $V$ .

If we have a splitting  $\mathcal{D}V = V \oplus V^*$ , then a generalized metric can be described in terms of forms. Indeed, in this case  $C_+$  can be described as a graph over  $V$  of an element in  $\text{Hom}(V, V^*) \cong \otimes^2 V^* = \text{Sym}^2 V^* \oplus \wedge^2 V^*$ , i.e., there is  $g \in \text{Sym}^2 V^*$  and  $b \in \wedge^2 V^*$  such that

$$C_+ = \{X + (b + g)(X) : X \in V\}.$$

It is clear that  $g$  above is nothing but the metric induced by  $\mathcal{G}$  on  $V$ . The subspace  $C_-$  is also a graph over  $V$ . Since it is orthogonal to  $C_+$  with respect to the natural pairing we see that

$$C_- = \{X + (b - g)(X) : X \in V\}.$$

Conversely, a metric  $g$  and a 2-form  $b$  define a pair of orthogonal spaces  $C_{\pm} \subset V \oplus V^*$  where the pairing is  $\pm$ -definite, so, on  $V \oplus V^*$ , a generalized metric is equivalent to a metric and a 2-form.

A generalized metric on  $V \oplus V^*$  allows us to define a Hodge star operator [35]:

**Definition 2.1.** Fix an orientation for  $C_+$  and let  $e_1, \dots, e_n$  be an oriented orthonormal basis for this space. Denoting by  $\tau$  the product  $e_1 \cdots e_n \in \text{Cl}(V \oplus V^*)$ , the *generalized Hodge star* is defined by  $\star \alpha = (-1)^{|\alpha|(n-1)} \tau \cdot \alpha$ , where  $\cdot$  is the Clifford action of  $\text{Cl}(V \oplus V^*)$  on forms.

If we denote by  $\star_g$  the usual Hodge star associated to the metric  $g$ , the Mukai pairing gives the following relation, if  $b = 0$ :

$$(\alpha, \star \beta) = \alpha \wedge \star_g \beta.$$

In the presence of a  $b$ -field, if we let  $\alpha = e^{-b}\tilde{\alpha}$  and  $\beta = e^{-b}\tilde{\beta}$ , then the relation becomes

$$(\alpha, \star\beta) = \tilde{\alpha} \wedge \star_g \tilde{\beta}.$$

Hence  $(\alpha, \star\alpha)$  is a nonvanishing volume form whenever  $\alpha \neq 0$ .

### 2.1.2 Hermitian structures

Given a generalized complex structure  $\mathcal{J}_1$  on a vector space  $V$ , we say a generalized metric  $\mathcal{G}$  is *compatible* with  $\mathcal{J}_1$  if they commute. In this situation,  $\mathcal{J}_2 = \mathcal{G}\mathcal{J}_1$  is automatically a generalized complex structure which commutes with  $\mathcal{G}$  and  $\mathcal{J}_1$ . Given a generalized complex structure on  $V$ , one can always find a generalized metric compatible with it.

If we let  $C_{\pm} \subset \mathcal{D}V$  be the  $\pm 1$ -eigenspaces of  $\mathcal{G}$ , then, using the fact that  $\mathcal{J}_1$  and  $\mathcal{G}$  commute, we see that  $\mathcal{J}_1 : C_{\pm} \rightarrow C_{\pm}$ . Since  $\pi : C_{\pm} \rightarrow V$  are isomorphisms, we can transport the complex structures on  $C_{\pm}$  to complex structures  $I_{\pm}$  on  $V$ . Furthermore, as  $\mathcal{J}_1$  is orthogonal with respect to the natural pairing,  $I_{\pm}$  are orthogonal with respect to the induced metric on  $V$ .

If the double of  $V$  is split as  $V \oplus V^*$ , then both  $\mathcal{J}_1$  and  $\mathcal{J}_2$  give rise to a decomposition of  $\wedge^{\bullet}V^* \otimes \mathbb{C}$  into their  $ik$ -eigenvalues. Since  $\mathcal{J}_1$  and  $\mathcal{J}_2$  commute, they can be diagonalized simultaneously:

$$U^{p,q} = U_{\mathcal{J}_1}^p \cap U_{\mathcal{J}_2}^q; \quad \oplus_{p,q} U^{p,q} = \wedge^{\bullet}V^* \otimes \mathbb{C}.$$

So, given a metric compatible with a generalized complex structure on  $V \oplus V^*$ , we obtain a  $\mathbb{Z}^2$  grading on forms. As we will see later, this bigrading will be compatible with the differentiable structure on a manifold only if the  $\mathcal{J}_i$  are integrable, which corresponds to the generalized Kähler case.

**Example 2.2.** Let  $V$  be a vector space endowed with a complex structure  $I$  compatible with a metric  $g$ . Then the induced generalized complex structure  $\mathcal{J}_I$  is compatible with the generalized metric  $\mathcal{G}$  on  $V \oplus V^*$  induced by  $g$  with  $b = 0$ . The generalized complex structure defined by  $\mathcal{G}\mathcal{J}_I$  is nothing but the generalized complex structure  $\mathcal{J}_{\omega}$  defined by the Kähler form  $\omega = g(I, \cdot)$ .



In this case, the decomposition of forms into  $U^{p,q}$  induced by this hermitian structure corresponds to the intersection of  $U_I^p$  and  $U_\omega^q$ , as determined in Examples 1.9 and 1.10, i.e.,

$$U^{p-q, n-p-q} = U_{\mathcal{J}_I}^{p-q} \cap U_{\mathcal{J}_\omega}^{n-p-q} = \Phi(\wedge^{p,q} V^*), \quad (2.1)$$

where  $\Phi(\alpha) = e^{i\omega} e^{\frac{\Delta}{2i}} \alpha$  is the map defined in Example 1.10. In particular, the decomposition of forms into  $U^{p,q}$  is not just the decomposition into  $\wedge^{p,q} V^*$ , which only depends on  $I$ , but isomorphic to it via an isomorphism which depends on the symplectic form.

Also, as we saw in the previous section, a metric on  $V \oplus V^*$  gives rise to a Hodge star operator. If the metric is compatible with a generalized complex structure, then this star operator can be expressed in terms of the Lie group action of  $\mathcal{J}_1$  and  $\mathcal{J}_2$  on forms:

**Lemma 2.3.** (Gualtieri [35]) *If a metric  $\mathcal{G}$  is compatible with a generalized complex structure  $\mathcal{J}_1$  and we let  $\mathcal{J}_2 = \mathcal{G}\mathcal{J}_1$ , then  $\star\alpha = \mathcal{J}_1\mathcal{J}_2\alpha$ .*

**Corollary 2.4.** *If  $V \oplus V^*$  has a generalized complex structure  $\mathcal{J}_1$  and a compatible metric  $\mathcal{G}$ , then the Hodge star operator preserves the bigrading of forms.*

*Proof.* Indeed if  $\alpha \in U^{p,q}$ , then  $\star\alpha = \mathcal{J}_1\mathcal{J}_2\alpha = i^{p+q}\alpha$ . □

For a generalized complex structure with compatible metric the operator  $\bar{\star}$  defined by  $\bar{\star}\alpha = \star\bar{\alpha}$  is also important, as it furnishes a definite, hermitian, bilinear functional,

$$h(\alpha, \beta) = (\alpha, \bar{\star}\beta), \quad \alpha, \beta \in \wedge^\bullet V^* \otimes \mathbb{C}. \quad (2.2)$$

We finish this section with a remark. The fact that a pair of commuting generalized complex structures  $\mathcal{J}_1$  and  $\mathcal{J}_2$  on  $V$  gives rise to a metric places algebraic restrictions on  $\mathcal{J}_1$  and  $\mathcal{J}_2$ . For example, one can check that,

$$\text{type}(\mathcal{J}_1) + \text{type}(\mathcal{J}_2) \leq n,$$

where  $\dim(V) = 2n$ .

## 2.2 Generalized Kähler structures

### 2.2.1 Hermitian structures

A Hermitian structure on an exact Courant algebroid  $\mathcal{E}$  over a manifold  $M$  is a generalized complex structure  $\mathcal{J}$  with compatible metric  $\mathcal{G}$  on  $\mathcal{E}$ . Given a generalized complex structure on  $\mathcal{E}$  one can always find a metric compatible with it, therefore obtaining a Hermitian structure. From a general point of view, this is true because it corresponds to a reduction of the structure group of the Courant algebroid from  $U(n, n)$  to its maximal compact subgroup  $U(n) \times U(n)$  and such reduction is unobstructed.

Incidentally, the existence of such a metric implies that every generalized almost complex manifold has an almost complex structure. Indeed, from the work in the previous section, if  $\mathcal{J}$  is a generalized almost complex structure, then a metric compatible with  $\mathcal{J}$  allows one to define an almost complex structure on  $M$  using the projection  $\pi : C_+ \rightarrow TM$ .

Given a splitting for  $\mathcal{E}$ ,  $\mathcal{J}$  induces a decomposition of differential forms,  $\mathcal{G}$  furnishes a Hodge star operator preserving this decomposition and the operator  $h$  from equation (2.2) is a definite hermitian bilinear functional.

**Lemma 2.5.** *Let  $(M, \mathcal{J}, \mathcal{G})$  be a generalized complex manifold with compatible metric. Then the  $h$ -adjoint of  $\bar{\partial}$  is given by  $\bar{\partial}^* = -\bar{\star}\bar{\partial}\bar{\star}^{-1}$*

*Proof.* We start observing that  $(d_H\alpha, \beta) + (\alpha, d_H\beta) = (d(\sigma(\alpha) \wedge \beta))_{top}$  is an exact form. Now, let  $\alpha \in \mathcal{U}^{k+1}$  and  $\beta \in \mathcal{U}^{-k}$ , then

$$\begin{aligned} (d(\sigma(\alpha) \wedge \beta))_{top} &= (d_H\alpha, \beta) + (\alpha, d_H\beta) \\ &= (\partial\alpha, \beta) + (\bar{\partial}\alpha, \beta) + (\alpha, \partial\beta) + (\alpha, \bar{\partial}\beta), \end{aligned} \tag{2.3}$$

and according to Proposition 1.8, the terms  $(\partial\alpha, \beta)$  and  $(\alpha, \partial\beta)$  vanish. Therefore

$$\begin{aligned}
h(\bar{\partial}\alpha, \beta) &= \int_M (\bar{\partial}\alpha, \star\bar{\beta}) = - \int_M (\alpha, \bar{\partial}\star\bar{\beta}) \\
&= - \int_M (\alpha, \overline{\star\star}^{-1}\bar{\partial}\star\bar{\beta}) \\
&= h(\alpha, -\overline{\star}^{-1}\bar{\partial}\overline{\star}\beta)
\end{aligned}$$

□

Now, the Laplacian  $\bar{\partial}\bar{\partial}^* + \bar{\partial}^*\bar{\partial}$  is an elliptic operator and in a compact generalized complex manifold every  $\bar{\partial}$ -cohomology class has a unique harmonic representative, which is a  $\bar{\partial}$  and  $\bar{\partial}^*$ -closed form. Also, from the expression above for  $\bar{\partial}^*$ , we see that  $\overline{\star}$  maps harmonic forms to harmonic forms.

**Theorem 2.6** (Serre duality; Cavalcanti [16]). *In a compact generalized complex manifold  $(M^{2n}, \mathcal{J})$ , the Mukai pairing gives rise to a pairing in cohomology  $H_{\bar{\partial}}^k \times H_{\bar{\partial}}^l \rightarrow H^{2n}(M)$  which vanishes if  $k \neq -l$  and is nondegenerate if  $k = -l$ .*

*Proof.* Given cohomology classes  $a \in H_{\bar{\partial}}^k(M)$  and  $b \in H_{\bar{\partial}}^l(M)$ , choose representative  $\alpha \in \mathcal{U}^k$  and  $\beta \in \mathcal{U}^l$ . According to Lemma 1.8,  $(\alpha, \beta)$  vanishes if  $k \neq -l$ , therefore proving the first claim.

If  $k = -l$  and  $b = 0$ , so that  $\beta = \bar{\partial}\gamma$  is a  $\bar{\partial}$ -exact form, then, according to (2.3),

$$[(\alpha, \bar{\partial}\gamma)] = [(\bar{\partial}\alpha, \gamma)] = 0,$$

Showing that the pairing is well defined.

Finally, if we let  $\alpha$  be the harmonic representative of the class  $a$ , then  $\overline{\star}\alpha$  is  $\bar{\partial}$  closed form in  $\mathcal{U}^{-k}$  which pairs nontrivially with  $\alpha$ , showing nondegeneracy. □

### 2.2.2 Generalized Kähler structures

Given a Hermitian structure  $(\mathcal{J}, \mathcal{G})$  on a Courant algebroid  $\mathcal{E}$ , we can always define the generalized almost complex structure  $\mathcal{J}_2 = \mathcal{J}\mathcal{G}$ . This structure is not integrable in general. Particular examples are

given by an almost complex structure taming a symplectic structure or by the Kähler form  $\omega = g(I, \cdot)$  on a Hermitian manifold. So the integrability of  $\mathcal{J}_2$  is the analogue of the Kähler condition.

**Definition 2.7.** A *generalized Kähler structure* on an exact Courant algebroid  $\mathcal{E}$  is a pair of commuting generalized complex structures  $\mathcal{J}_1$  and  $\mathcal{J}_2$  such that  $\mathcal{G} = -\mathcal{J}_1\mathcal{J}_2$  is a generalized metric.

As we have seen, given a generalized complex structure compatible with a generalized metric one can define almost complex structures  $I_{\pm}$  on  $TM$  using the projections  $\pi : C_{\pm} \rightarrow TM$ . The integrability of  $\mathcal{J}_1$  and  $\mathcal{J}_2$  imply that  $I_{\pm}$  are integrable complex structures and also that

$$dd^c_{-}\omega_{-} = 0 \quad d^c_{-}\omega_{-} = -d^c_{+}\omega_{+},$$

where  $\omega_{\pm} = g(I_{\pm}, \cdot)$  are the Kähler forms associated to  $I_{\pm}$  and  $d^c_{\pm} = -I_{\pm}dI_{\pm}$  (see [34], Proposition 6.16).

The converse also holds and is the heart of Gualtieri's theorem relating generalized Kähler structure to bihermitian structures:

**Theorem 2.8** (Gualtieri [34], Theorem 6.37). *A manifold  $M$  admits a bihermitian structure  $(g, I_{+}, I_{-})$ , such that*

$$dd^c_{-}\omega_{-} = 0 \quad d^c_{-}\omega_{-} = -d^c_{+}\omega_{+}$$

*if and only if the Courant algebroid  $\mathcal{E}$  with characteristic class  $[d^c_{-}\omega_{-}]$  has a generalized Kähler structure which induces the bihermitian structure  $(g, I_{+}, I_{-})$  on  $M$ .*

In the case of a generalized Kähler induced by a Kähler structure,  $I_{+} = \pm I_{-}$  and the equations above hold trivially as  $d^c\omega = 0$ . Now we use the bihermitian characterization of a generalized Kähler manifold to give nontrivial examples of generalized Kähler structures.

**Example 2.9** (Gualtieri [34], Example 6.30). Let  $(M, g, I, J, K)$  be a hyperkähler manifold. Then it automatically has an  $S^2$  worth of Kähler structures which automatically furnish generalized Kähler structures. However there are other generalized Kähler structures on  $M$ . For example, if we let  $I_{+} = I$  and  $I_{-} = J$ , then all the conditions of Theorem 2.8 hold hence providing  $M$  with a generalized Kähler

structure (with  $H = 0$ ). The forms generating the canonical bundles of this generalized Kähler structure are

$$\rho_1 = \exp(\omega_K + \frac{i}{2}(\omega_I - \omega_J));$$

$$\rho_2 = \exp(-\omega_K + \frac{i}{2}(\omega_I + \omega_J)).$$

showing that a generalized Kähler structure can be determined by two generalized complex structures of symplectic type.

**Example 2.10** (Gualtieri [34], Example 6.39). Every compact even dimensional Lie group  $G$  admits left and right invariant complex structures [58, 64]. If  $G$  is semi-simple, we can choose such complex structures to be Hermitian with respect to the invariant metric induced by the Killing form  $K$ . This bihermitian structure furnishes a generalized Kähler structure on the Courant algebroid over  $G$  with characteristic class given by the bi-invariant Cartan 3-form:  $H(X, Y, Z) = K([X, Y], Z)$ . To prove this we let  $J_L$  and  $J_R$  be left and right invariant complex structures as above and compute  $d_L^c \omega_L$ :

$$\begin{aligned} A &= d_L^c \omega_L(X, Y, Z) = d_L \omega_L(J_L X, J_L Y, J_L Z) \\ &= -\omega_L([J_L X, J_L Y], J_L Z) + c.p. \\ &= -K([J_L X, J_L Y], Z) + c.p. \\ &= -K(J_L[J_L X, Y] + J_L[X, J_L Y] + [X, Y], Z) + c.p. \\ &= (2K([J_L X, J_L Y], Z) + c.p.) - 3H(X, Y, Z) \\ &= -2A - 3H, \end{aligned}$$

where  $c.p.$  stands for cyclic permutations. This proves that  $d_L^c \omega_L = -H$ . Since the right Lie algebra is antiholomorphic to the left, the same calculation yields  $d_R^c \omega_R = H$  and by Theorem 2.8, this bihermitian structure induces a generalized Kähler structure on the Courant algebroid with characteristic class  $[H]$ .

More recently, using a classification theorem for bihermitian structures on 4-manifolds by Apostolov, Gauduchon and Grantcharov [1], Apostolov and Gualtieri managed to classify all 4-manifolds admitting generalized Kähler structures [2]. Obviously, all Kähler 4-manifolds are on that list. The question of which Kähler 4-manifolds admit non-Kähler generalized Kähler structures is still open.

## 2.3 Hodge identities

Assume that an exact Courant algebroid  $\mathcal{E}$  has a generalized Kähler structure  $(\mathcal{J}_1, \mathcal{J}_2)$ . Once a splitting for  $\mathcal{E}$  is chosen, we obtain a bigrading of forms into  $U^{p,q} = U_{\mathcal{J}_1}^p \cap U_{\mathcal{J}_2}^q$ . Since for generalized complex structure  $d_H : \mathcal{U}^k \longrightarrow \mathcal{U}^{k+1} + \mathcal{U}^{k-1}$ , we see that for a generalized Kähler structure  $d_H$  decomposes in four components:

$$d_H : \mathcal{U}^{p,q} \longrightarrow \mathcal{U}^{p+1,q+1} + \mathcal{U}^{p+1,q-1} + \mathcal{U}^{p-1,q+1} + \mathcal{U}^{p-1,q-1}.$$

We denote these components by

$$\delta_+ : \mathcal{U}^{p,q} \longrightarrow \mathcal{U}^{p+1,q+1} \quad \delta_- : \mathcal{U}^{p,q} \longrightarrow \mathcal{U}^{p+1,q-1}$$

and their conjugates

$$\overline{\delta}_+ : \mathcal{U}^{p,q} \longrightarrow \mathcal{U}^{p-1,q-1} \quad \overline{\delta}_- : \mathcal{U}^{p,q} \longrightarrow \mathcal{U}^{p-1,q+1}.$$

So that, for example,  $\partial_{\mathcal{J}_1} = \delta_+ + \delta_-$  and  $\partial_{\mathcal{J}_2} = \delta_+ + \overline{\delta}_-$ . One can easily show that  $h$ -adjoints of  $\delta_{\pm}$  are given by  $\delta_{\pm}^* = -\overline{\delta}_{\pm} \star^{-1}$  and similarly for  $\overline{\delta}_-$ ,  $\overline{\delta}_+$  and  $\delta_-$ . Using the description of  $\star$  in terms of the Lie algebra action of  $\mathcal{J}_1$  and  $\mathcal{J}_2$  given in Lemma 2.3, Gualtieri proved

**Theorem 2.11** (Gualtieri [35]). *The following relations hold in a generalized Kähler manifold*

$$\delta_+^* = \overline{\delta}_+ \quad \text{and} \quad \delta_-^* = -\overline{\delta}_-; \quad (2.4)$$

$$\begin{aligned} 4\Delta_{d_H} &= 2\Delta_{\partial_{\mathcal{J}_1}} = 2\Delta_{\overline{\delta}_+} = 2\Delta_{\partial_{\mathcal{J}_2}} = 2\Delta_{\overline{\delta}_-} \\ &= \Delta_{\delta_+} + \Delta_{\delta_-} = \Delta_{\overline{\delta}_+} + \Delta_{\overline{\delta}_-}. \end{aligned} \quad (2.5)$$

*Proof.* To prove that  $\delta_+^* = -\overline{\delta}_+$ , one only has to prove these operators agree when action on  $\mathcal{U}^{p,q}$ . So let  $\alpha \in \mathcal{U}^{p,q}$ . Then, according to Lemma 2.3, we get  $\overline{\delta}_+ \alpha = i^{-p-q} \overline{\alpha}$  and

$$\delta_+^* \alpha = -\overline{\delta}_+ \star^{-1} \alpha = -i^{p+q} \overline{\delta}_+ \overline{\alpha} = -i^{p+q} i^{-p+1-q+1} \overline{\delta}_+ \overline{\alpha} = \overline{\delta}_+ \alpha$$

And similarly one can check the identity  $\delta_-^* = -\overline{\delta}_-$ .

To prove the equality of the Laplacians now one only has to observe that due to (2.4)

$$\Delta_{\delta_+} = \delta_+ \bar{\delta}_+ + \bar{\delta}_+ \delta_+ \text{ and } \Delta_{\delta_-} = -(\delta_- \bar{\delta}_- + \bar{\delta}_- \delta_-).$$

On the other hand, if  $\alpha \in \mathcal{U}^{p,q}$  then  $d_H^2 \alpha = 0$  implies

$$(\delta_+ \bar{\delta}_+ + \bar{\delta}_+ \delta_+ + \delta_- \bar{\delta}_- + \bar{\delta}_- \delta_-) \alpha = 0$$

for all  $\alpha$  and hence  $\Delta_{\delta_+} = \Delta_{\delta_-}$ .

Since  $\partial_{\mathcal{J}_1}^2 = 0$ ,  $\delta_+$  and  $\delta_-$  anticommute, so, e.g.,

$$\begin{aligned} \Delta_{d_H} &= d_H d_H^* + d_H^* d_H = (\delta_+ + \delta_- + \bar{\delta}_+ + \bar{\delta}_-)(\delta_+^* + \delta_-^* + \bar{\delta}_+^* + \bar{\delta}_-^*) \\ &\quad + (\delta_+^* + \delta_-^* + \bar{\delta}_+^* + \bar{\delta}_-^*)(\delta_+ + \delta_- + \bar{\delta}_+ + \bar{\delta}_-) \\ &= (\delta_+ + \delta_- + \bar{\delta}_+ + \bar{\delta}_-)(\bar{\delta}_+ - \bar{\delta}_- + \delta_+ - \delta_-) \\ &\quad + (\bar{\delta}_+ - \bar{\delta}_- + \delta_+ - \delta_-)(\delta_+ + \delta_- + \bar{\delta}_+ + \bar{\delta}_-) \\ &= 2(\delta_+ \bar{\delta}_+ + \bar{\delta}_+ \delta_+) + 2(\delta_- \bar{\delta}_- + \bar{\delta}_- \delta_-) \\ &= 2\Delta_{\delta_+} + 2\Delta_{\delta_-} = 4\Delta_{\delta_+}. \end{aligned}$$

And similar arguments prove all the other identities.  $\square$

Here we mention a couple of standard consequences of this theorem whose proof follows the same argument given in the classical Kähler case.

**Corollary 2.12** (Gualtieri [35]). *In a compact generalized Kähler manifold the decomposition of forms into  $\mathcal{U}^{p,q}$  gives rise to a  $(p, q)$ -decomposition of the  $d_H$ -cohomology.*

**Corollary 2.13** (Gualtieri [35]).  **$(\delta_+ \delta_-$ -Lemma)** *In a compact generalized Kähler manifold*

$$\text{Im}(\delta_+) \cap \text{Ker}(\delta_-) = \text{Im}(\delta_-) \cap \text{Ker}(\delta_+) = \text{Im}(\delta_+ \delta_-).$$

## 2.4 Formality in generalized Kähler geometry

A differential graded algebra  $(\mathcal{A}, d)$  is *formal* if there is a finite sequence of differential graded algebras  $(\mathcal{A}_i, d)$  with quasi isomorphisms

$\varphi_i$  between  $\mathcal{A}_i$  and  $\mathcal{A}_{i+1}$  such that  $(\mathcal{A}_1, d) = (\mathcal{A}, d)$  and  $(\mathcal{A}_n, d) = (H^\bullet(A), 0)$ :

$$\mathcal{A}_1 \xrightarrow{\varphi_1} \mathcal{A}_2 \xleftarrow{\varphi_2} \mathcal{A}_3 \xrightarrow{\varphi_3} \dots \xrightarrow{\varphi_{n-2}} \mathcal{A}_{n-1} \xleftarrow{\varphi_{n-1}} \mathcal{A}_n.$$

And a manifold is *formal* if the algebra of differential forms,  $(\Omega^\bullet(M), d)$ , is formal.

One of the most striking uses of the  $\partial\bar{\partial}$ -lemma for a complex structure appears in the proof that it implies formality [25], therefore providing fine topological obstructions for a manifold to admit a Kähler structure [61]. A very important fact used in this proof is that for a complex structure  $\partial$  and  $\bar{\partial}$  are derivations. As we mentioned before, in the generalized complex world the operators  $\partial$  and  $\bar{\partial}$  are not derivations and indeed there are examples of nonformal generalized complex manifolds for which the  $\partial\bar{\partial}$ -lemma holds [17].

However, as we saw in Section 1.3.1, given a generalized complex structure we can form the differential graded algebra  $(\Omega^\bullet(\bar{L}), d_L)$  and if the generalized complex structure has holomorphically trivial canonical bundle, trivialized by a form  $\rho$ , we get a map

$$\phi : \Omega^\bullet(\bar{L}) \longrightarrow \Omega_{\mathbb{C}}^\bullet(M); \quad \phi(s \cdot \sigma) = s \cdot \rho$$

such that

$$\bar{\partial}\phi(\alpha) = \phi(d_L\alpha).$$

In the case of a generalized Kähler structure, since  $\mathcal{J}_1$  and  $\mathcal{J}_2$  commute,  $\mathcal{J}_2$  furnishes an integrable complex structure on  $L_1$ , the  $i$ -eigenspace of  $\mathcal{J}_1$ . With that we obtain a  $p, q$ -decomposition of  $\wedge^\bullet \bar{L}_1$  and decomposition  $d_{L_1} = \partial_{L_1} + \bar{\partial}_{L_1}$  of the differential. Since this is nothing but the decomposition of  $d_{L_1}$  by an underlying complex structure, the operators  $\partial_{L_1}$  and  $\bar{\partial}_{L_1}$  are derivations. And if further  $\mathcal{J}_1$  has holomorphically trivial canonical bundle, then the correspondence between  $d_L$  and  $\partial$  gives an identifications between  $\partial_{L_1}$  and  $\delta_+$  and  $\bar{\partial}_{L_1}$  and  $\delta_-$ :

$$\delta_+\phi(\alpha) = \phi(\partial_{L_1}\alpha) \quad \delta_-\phi(\alpha) = \phi(\bar{\partial}_{L_1}\alpha).$$

So, if we let  $d_{L_1}^c = -i(\partial_{L_1} - \bar{\partial}_{L_1})$ , Corollary 2.13 implies



**Lemma 2.14.** *If  $(\mathcal{J}_1, \mathcal{J}_2)$  is a generalized Kähler structure on a compact manifold and  $\mathcal{J}_1$  has holomorphically trivial canonical bundle then*

$$\mathrm{Im}(d_{L_1}) \cap \mathrm{Ker}(d_{L_1}^c) = \mathrm{Im}(d_{L_1}^c) \cap \mathrm{Ker}(d_{L_1}) = \mathrm{Im}(d_{L_1} d_{L_1}^c).$$

And hence the same argument from [25] gives

**Theorem 2.15** (Cavalcanti [15]). *If  $(\mathcal{J}_1, \mathcal{J}_2)$  is a generalized Kähler structure on a compact manifold and  $\mathcal{J}_1$  has holomorphically trivial canonical bundle, then  $(\Omega^\bullet(\overline{L}_1), d_{L_1})$  is a formal differential graded algebra.*

*Proof.* Let  $(\Omega_c^\bullet(\overline{L}_1), d_{L_1})$  be the algebra of  $d_{L_1}^c$ -closed element of  $\Omega^\bullet(\overline{L}_1)$  endowed with differential  $d_{L_1}$  and  $(H_{d_{L_1}^c}(\overline{L}_1), d_{L_1})$  be the cohomology of  $\Omega^\bullet(\overline{L}_1)$  with respect to  $d_{L_1}^c$ , also endowed with differential  $d_{L_1}$ . Then we have maps

$$(\Omega(\overline{L}_1), d_{L_1}) \xleftarrow{i} (\Omega_c(\overline{L}_1), d_{L_1}) \xrightarrow{\pi} (H_{d_{L_1}^c}(\overline{L}_1), d_{L_1}),$$

and, as we are going to see, these maps are quasi-isomorphisms and the differential of  $(H_{d_{L_1}^c}(M), d_{L_1})$  is zero, therefore showing that  $\Omega(\overline{L}_1)$  is formal.

i)  $i^*$  is surjective:

Given a  $d_{L_1}$ -closed form  $\alpha$ , let  $\beta = d_{L_1}^c \alpha$ . Then  $d_{L_1} \beta = d_{L_1} d_{L_1}^c \alpha = -d_{L_1}^c d_{L_1} \alpha = 0$ , so  $\beta$  satisfies the conditions of the  $d_{L_1} d_{L_1}^c$ -lemma, hence  $\beta = d_{L_1}^c d_{L_1} \gamma$ . Let  $\tilde{\alpha} = \alpha - d_{L_1} \gamma$ , then  $d_{L_1}^c \tilde{\alpha} = d_{L_1}^c \alpha - d_{L_1}^c d_{L_1} \gamma = \beta - \beta = 0$ , so  $[\alpha] \in \mathrm{Im}(i^*)$ .

ii)  $i^*$  is injective:

If  $i^* \alpha$  is exact, then  $\alpha$  is  $d_{L_1}^c$ -closed and exact, hence by the  $d_{L_1} d_{L_1}^c$ -lemma  $\alpha = d_{L_1} d_{L_1}^c \beta$ , so  $\alpha$  is the derivative of a  $d_{L_1}^c$ -closed form and hence its cohomology class in  $\Omega_c$  is also zero.

iii) The differential of  $(H_{d_{L_1}^c}(M), d)$  is zero:

Let  $\alpha$  be  $d_{L_1}^c$ -closed, then  $d_{L_1} \alpha$  is exact and  $d_{L_1}^c$  closed so, by the  $d_{L_1} d_{L_1}^c$ -lemma,  $d_{L_1} \alpha = d_{L_1}^c d_{L_1} \beta$  and so it is zero in  $d_{L_1}^c$ -cohomology.

iv)  $\pi^*$  is onto:

Let  $\alpha$  be  $d_{L_1}^c$ -closed. Then, as above,  $d_{L_1}\alpha = d_{L_1}d_{L_1}^c\beta$ . Let  $\tilde{\alpha} = \alpha - d_{L_1}^c\beta$ , and so  $d_{L_1}\tilde{\alpha} = 0$  and  $[\tilde{\alpha}]_{d_{L_1}^c} = [\alpha]_{d_{L_1}^c}$ , so  $\pi^*([\tilde{\alpha}]_{d_{L_1}}) = [\alpha]_{d_{L_1}^c}$ .

v)  $\pi^*$  is injective:

Let  $\alpha$  be closed and  $d_{L_1}^c$ -exact, then the  $d_{L_1}d_{L_1}^c$ -lemma implies that  $\alpha$  is exact and hence  $[\alpha] = 0$  in  $\mathcal{E}_c$ .

□

If  $\mathcal{J}_1$  is a structure of type 0, i.e., is of symplectic type, then not only does it have holomorphically trivial canonical bundle but  $\pi : L \rightarrow T_{\mathbb{C}}M$  is an isomorphism and the bracket on  $L$  is mapped to the Lie bracket of vector fields. Therefore, in this case,  $(\Omega(\overline{L}_1), d_{L_1})$  is isomorphic to  $(\Omega_{\mathbb{C}}(M), d)$ . So the previous theorem gives:

**Corollary 2.16** (Cavalcanti [15]). *If  $(\mathcal{J}_1, \mathcal{J}_2)$  is a generalized Kähler structure on a compact manifold  $M$  and  $\mathcal{J}_1$  is of symplectic type, then  $M$  is formal.*

Similarly to the original theorem of formality of Kähler manifolds, Theorem 2.15 furnishes a nontrivial obstruction for a given generalized complex structure to be part of a generalized Kähler structure. As an application of this result one can prove that no generalized complex structure on a nilpotent Lie algebra is part of generalized Kähler pair [15].

## Chapter 3

# Reduction of Courant algebroids and generalized complex structures

Given a structure on a manifold  $M$  and a group  $G$  acting on  $M$  by symmetries of that structure, one can ask what kind of conditions have to be imposed on the group action in order for that structure on  $M$  to descend to a similar type of structure on  $M/G$ . Examples include the quotient of metrics when a Riemannian manifold is acted on by Killing fields, quotient of complex manifolds by holomorphic actions of a complex group and the *reduction* of symplectic manifolds acted on by a group of symplectomorphisms.

The latter example is particularly interesting as it shows that sometimes it may be necessary to take submanifolds as well as the quotient by the  $G$  action in order to find a manifold with the desired structure. This is going to be a central feature in theory developed in this chapter and one of the tasks ahead is to define a notion of action which includes the choice of submanifolds on it. Although we are primarily concerned about how to take quotients of generalized

complex structures, there is a more basic question which needs to be answered first: how can we quotient a Courant algebroid  $\mathcal{E}$ ?

To answer this question we recall that the action of  $G$  on  $M$  can be fully described by the infinitesimal action of its Lie algebra,  $\psi : \mathfrak{g} \rightarrow \Gamma(TM)$ , which is a morphism of Lie algebras. We want to have a similar picture for a Courant algebroid  $\mathcal{E}$  over  $M$ , i.e., we want to describe a  $G$ -action on  $\mathcal{E}$  covering the  $G$ -action on  $M$  by a map  $\Psi : \mathfrak{a} \rightarrow \Gamma(\mathcal{E})$ . Here we encounter our first problem which is to determine what kind of object  $\mathfrak{a}$  is. It is natural to ask that it has the same sort of structure that  $\mathcal{E}$  has, i.e., instead of being a Lie algebra, it has to encode the properties of a Courant algebroid. This leads us to the concept of a *Courant algebra*. Another point is that we still want to have the same group  $G$  acting on both  $\mathcal{E}$  and  $M$ , thus restricting the Courant algebra morphisms  $\Psi$  one is allowed to consider: these are the *extended actions*.

We will see that once an extended action on a Courant algebroid is chosen, it determines a foliation on  $M$  whose leaves are invariant under the  $G$  action. The ‘quotient’ algebroid is an algebroid defined over the quotient of a leaf of this distribution by  $G$  and the algebroid itself is obtained by considering the quotient of a subspace of  $\mathcal{E}$  and hence we call these the *reduced manifold* and the *reduced Courant algebroid*. Some of the formalism we introduce appeared before in the physics literature in the context of gauging the Wess–Zumino term in a sigma model [40, 29, 28].

Once the reduced Courant algebroid  $\mathcal{E}_{red}$  is understood, it is relatively easy to reduce structures from  $\mathcal{E}$ . Here we will deal only with Dirac and generalized complex structures. However, even by just considering these we see that this theory is actually quite strong and includes as subcases pull back and push forward of Dirac structures (as studied in [12]), quotient of complex manifolds by holomorphic actions by complex groups and symplectic reduction. Other interesting and new cases consist of type changing reductions: for example, the reduction of a symplectic structure can lead to a generalized complex structure of nonzero type (see Example 3.24).

The material of this chapter is mostly an extraction from the collaboration with Bursztyn and Gualtieri [11], whose reading I highly recommend if you are interested in these topics. There we also deal with the reduction of generalized Kähler structures and present the

concept of Hamiltonian actions. Other independent work dealing with reduction of generalized complex structures with different degrees of generality are [38, 48, 60].

This chapter is organized as follows. In Section 3.1 we give a precise description of the group of symmetries of an exact Courant algebroid, define Courant algebras and extended actions. In Section 3.2 given an extended action on an exact Courant algebroid over a manifold, we explain how to reduce the manifold and the Courant algebroid. In Section 3.3 we show how to transport Dirac structures from the original Courant algebroid to the reduced Courant algebroid and then we finish in Section 3.4 applying these results to reduction of generalized complex structures.

## 3.1 Courant algebras and extended actions

### 3.1.1 Symmetries of Courant algebroids

As we have mentioned before, the group of symmetries,  $\mathcal{C}$ , of an exact Courant algebroid is formed by diffeomorphisms and  $B$ -fields. A more precise description of  $\mathcal{C}$  can be given once an isotropic splitting is chosen [34]: it consists of the group of ordered pairs  $(\varphi, B) \in \text{Diff}(M) \times \Omega^2(M)$  such that  $\varphi^*H - H = dB$ , where  $H$  is the curvature of the splitting. Diffeomorphisms act in the usual way on  $TM \oplus T^*M$ , while 2-forms act via  $B$ -field transforms. As a result we see that  $\mathcal{C}$  is an extension

$$0 \longrightarrow \Omega_{cl}^2(M) \longrightarrow \mathcal{C} \longrightarrow \text{Diff}_{[H]}(M) \longrightarrow 0,$$

where  $\text{Diff}_{[H]}(M)$  is the group of diffeomorphisms preserving the cohomology class  $[H]$ .

Therefore, the Lie algebra  $\mathfrak{c}$  of symmetries consists of pairs  $(X, B) \in \Gamma(TM) \oplus \Omega^2(M)$  such that  $\mathcal{L}_X H = dB$ . For this reason, it is an extension of the form

$$0 \longrightarrow \Omega_{cl}^2(M) \longrightarrow \mathfrak{c} \longrightarrow \Gamma(T) \longrightarrow 0.$$

We have mentioned before that there is a natural adjoint action of a section  $e_1$  of  $\mathcal{E}$  on  $\Gamma(\mathcal{E})$  by  $e_1 \bullet e_2 = \llbracket e_1, e_2 \rrbracket + d\langle e_1, e_2 \rangle$ . It follows

from the definition of a Courant algebroid that the adjoint action of  $e_1$  is an infinitesimal symmetry of  $\mathcal{E}$ , i.e.,

$$\pi(e_1)\langle e_2, e_3 \rangle = \langle e_1 \bullet e_2, e_3 \rangle + \langle e_2, e_1 \bullet e_3 \rangle \quad (3.1)$$

$$e_1 \bullet \llbracket e_2, e_3 \rrbracket = \llbracket e_1 \bullet e_2, e_3 \rrbracket + \llbracket e_2, e_1 \bullet e_3 \rrbracket, \quad (3.2)$$

and hence we have a map  $\text{ad} : \Gamma(\mathcal{E}) \rightarrow \mathfrak{c}$ . However, unlike the usual adjoint action of vector fields on the tangent bundle,  $\text{ad}$  is neither surjective nor injective; instead, in an exact Courant algebroid, the Lie algebra  $\mathfrak{c}$  fits into the following exact sequence:

$$0 \longrightarrow \Omega_{cl}^1(M) \xrightarrow{\frac{1}{2}\pi^*} \Gamma(\mathcal{E}) \longrightarrow \mathfrak{c} \longrightarrow H^2(M, \mathbb{R}) \longrightarrow 0,$$

where the map to cohomology can be written as  $(X, B) \mapsto [i_X H - B]$  in a given splitting.

### 3.1.2 Extended actions

Similarly to the case of a Lie group action, an infinitesimal action on an exact Courant algebroid is described in terms of sections of  $\mathcal{E}$ , using the argument above to identify them with symmetries. However, since the Courant bracket in  $\Gamma(\mathcal{E})$  is not a Lie bracket, the action is not given by a Lie algebra homomorphism, but by a map of *Courant algebras* (see definition below), an algebraic structure designed to capture the information regarding the Courant bracket and the pairing on  $\mathcal{E}$ .

**Definition 3.1.** A *Courant algebra* over the Lie algebra  $\mathfrak{g}$  is a vector space  $\mathfrak{a}$  equipped with a skew-symmetric bracket  $\llbracket \cdot, \cdot \rrbracket : \mathfrak{a} \times \mathfrak{a} \rightarrow \mathfrak{a}$ , a symmetric bilinear operation  $\theta : \mathfrak{a} \times \mathfrak{a} \rightarrow \mathfrak{a}$ , and a map  $\pi : \mathfrak{a} \rightarrow \mathfrak{g}$ , which satisfy the following conditions for all  $a_1, a_2, a_3 \in \mathfrak{a}$ :

- c1)  $\pi(\llbracket a_1, a_2 \rrbracket) = [\pi(a_1), \pi(a_2)]$ ,
- c2)  $\text{Jac}(a_1, a_2, a_3) = \frac{1}{3}(\theta(\llbracket a_1, a_2 \rrbracket), a_3) + c.p.)$ ,
- c3)  $\theta(a_1, a_2) \bullet a_3 = 0$ ,
- c4)  $\pi \circ \theta = 0$ ,

$$\text{c5) } a_1 \bullet \theta(a_2, a_3) = \theta(a_1 \bullet a_2, a_3) + \theta(a_2, a_1 \bullet a_3),$$

where  $\bullet$  denotes the combination

$$a_1 \bullet a_2 = \llbracket a_1, a_2 \rrbracket + \theta(a_1, a_2). \quad (3.3)$$

A Courant algebra is *exact* if  $\pi : \mathfrak{a} \rightarrow \mathfrak{g}$  is onto and  $\llbracket h_1, h_2 \rrbracket = \theta(h_1, h_2) = 0$  for all  $h_i \in \mathfrak{h} = \ker \pi$ .

One can ditch the skew bracket  $\llbracket \cdot, \cdot \rrbracket$  and the bilinear form  $\theta$  above and instead work with the nonskew bracket  $\bullet$ . From this angle, equation (3.3) shows how to define  $\llbracket \cdot, \cdot \rrbracket$  and  $\theta$  as the skew and symmetric parts of  $\bullet$ .

**Exercise 3.2.** A Courant algebra can alternatively be defined as a vector space  $\mathfrak{a}$  together with a bracket  $\bullet$  satisfying the Leibniz rule

$$a_1 \bullet (a_2 \bullet a_3) = (a_1 \bullet a_2) \bullet a_3 + a_2 \bullet (a_1 \bullet a_3),$$

and a map to Lie algebra  $\mathfrak{g}$ ,  $\pi : \mathfrak{a} \rightarrow \mathfrak{g}$ , which preserves brackets

$$\pi(a_1 \bullet a_2) = [\pi(a_1), \pi(a_2)].$$

An exact Courant algebra is one for which  $\pi$  is surjective and  $\bullet$  vanishes on  $\mathfrak{h} \otimes \mathfrak{h}$ .

A Courant algebroid  $\mathcal{E}$  gives an example of a Courant algebra over  $\mathfrak{g} = \Gamma(TM)$ , taking  $\mathfrak{a} = \Gamma(\mathcal{E})$  and  $\theta(e_1, e_2) = d\langle e_1, e_2 \rangle$ .  $\mathcal{E}$  is an exact Courant algebroid if and only if  $\Gamma(\mathcal{E})$  is an exact Courant algebra.

Given an exact Courant algebra, one obtains immediately an action of  $\mathfrak{g}$  on  $\mathfrak{h}$ :  $g \in \mathfrak{g}$  acts on  $h \in \mathfrak{h}$  via  $g \cdot h = a \bullet h$ , for any  $a$  such that  $\pi(a) = g$ . This is well defined, since  $\bullet$  vanishes on  $\mathfrak{h} \times \mathfrak{h}$ , and it determines an action because of the Leibniz property of  $\bullet$ .

$$\begin{aligned} g_1 \cdot (g_2 \cdot h) - g_2 \cdot (g_1 \cdot h) &= a_1 \bullet (a_2 \bullet h) - a_2 \bullet (a_1 \bullet h) \\ &= (a_1 \bullet a_2) \bullet h + a_2 \bullet (a_1 \bullet h) - a_2 \bullet (a_1 \bullet h) \\ &= (\llbracket a_1, a_2 \rrbracket + \theta(a_1, a_2)) \bullet h \\ &= \llbracket a_1, a_2 \rrbracket \bullet h = [g_1, g_2] \cdot h \quad \forall h \in \mathfrak{h}. \end{aligned}$$

A partial converse to this fact is given by the following example.

**Example 3.3** (Demisemidirect product [11, 43]). Let  $\mathfrak{g}$  be a Lie algebra acting on the vector space  $\mathfrak{h}$ . Then the following bracket and bilinear form make  $\mathfrak{a} = \mathfrak{g} \oplus \mathfrak{h}$  a Courant algebra over  $\mathfrak{g}$

$$\begin{aligned} \llbracket (g_1, h_1), (g_2, h_2) \rrbracket &= ([g_1, g_2], \tfrac{1}{2}(g_1 \cdot h_2 - g_2 \cdot h_1)), \\ \theta((g_1, h_1), (g_2, h_2)) &= (0, \tfrac{1}{2}(g_1 \cdot h_2 + g_2 \cdot h_1)), \end{aligned}$$

where here  $g \cdot h$  denotes the  $\mathfrak{g}$ -action. Indeed, conditions c1) and c4) are evident. Condition c2) follows from the Leibniz rule for  $\mathfrak{g}$  and the fact the  $\mathfrak{h}$  is a  $\mathfrak{g}$ -module:

$$\begin{aligned} \text{Jac}((g_1, h_1), (g_2, h_2), (g_3, h_3)) &= \\ &= (\text{Jac}(g_1, g_2, g_3), \tfrac{1}{2}([g_1, g_2] \cdot h_3 - g_3 \cdot \tfrac{1}{2}(g_1 \cdot h_2 - g_2 \cdot h_1)) + c.p.) \\ &= (0, \tfrac{1}{4}([g_1, g_2] \cdot h_3 + c.p.)) \\ &= \tfrac{1}{3}\theta((g_1, h_1), (g_2, h_2), (g_3, h_3)). \end{aligned}$$

Conditions c3) and c5) can be easily checked once we write the expression for the adjoint action

$$(g_1, h_1) \bullet (g_2, h_2) = ([g_1, g_2], g_1 \cdot h_2). \quad (3.4)$$

Since  $\theta(a_1, a_2) = (0, h)$ , for some  $h \in \mathfrak{h}$ , the expression above shows that c3) holds and c5) is not difficult to check either.

This bracket has appeared before in the context of Leibniz algebras [43], where it was called the *demisemidirect* product, due to the factor of  $\frac{1}{2}$ . Note that in [66], Weinstein studied the case where  $\mathfrak{g} = \mathfrak{gl}(V)$  and  $\mathfrak{h} = V$ , and called it an *omni-Lie algebra* due to the fact that, when  $\dim V = n$ , any  $n$ -dimensional Lie algebra can be embedded inside  $\mathfrak{g} \oplus \mathfrak{h}$  as an involutive subspace.

**Exercise 3.4.** Check that the demisemidirect product is a Courant algebra using the definition from Exercise 3.2.

Definition 3.1 is a pedagogical one, as it makes clear the analogies between the definitions of a Courant algebra and a Courant algebroid. However, solving the previous exercise shows that Definition 3.2 is much more treatable (for one thing, it has less conditions to check). Therefore, often in the sequel we will use 3.2 as the definition of Courant algebra.



**Definition 3.5.** A morphism of Courant algebras from  $(\mathfrak{a} \xrightarrow{\pi} \mathfrak{g}, [\cdot, \cdot], \theta)$  to  $(\mathfrak{a}' \xrightarrow{\pi'} \mathfrak{g}', [\cdot, \cdot]', \theta')$  is a commutative square

$$\begin{array}{ccc} \mathfrak{a} & \xrightarrow{\pi} & \mathfrak{g} \\ \Psi \downarrow & & \downarrow \psi \\ \mathfrak{a}' & \xrightarrow{\pi'} & \mathfrak{g}' \end{array}$$

where  $\psi$  is a Lie algebra homomorphism,  $\Psi(\llbracket a_1, a_2 \rrbracket) = \llbracket \Psi(a_1), \Psi(a_2) \rrbracket'$  and  $\Psi(\theta(a_1, a_2)) = \theta'(\Psi(a_1), \Psi(a_2))$  for all  $a_i \in \mathfrak{a}$ . Note that a morphism of Courant algebras induces a chain homomorphism of associated chain complexes  $\mathfrak{h} \longrightarrow \mathfrak{a} \xrightarrow{\pi} \mathfrak{g}$ .

Given a Courant algebra morphism  $\Psi : \mathfrak{a} \longrightarrow \Gamma(\mathcal{E})$  we can compose  $\Psi$  with  $\Phi : \Gamma(\mathcal{E}) \longrightarrow \mathfrak{c}$  to obtain a subalgebra of the algebra of infinitesimal symmetries of  $\mathcal{E}$ . Also, projecting  $\Psi$  onto  $\Gamma(TM)$  we obtain a subalgebra of infinitesimal symmetries of  $M$ .

$$\begin{array}{ccccc} \mathfrak{a} & \xrightarrow{\Psi} & \Gamma(\mathcal{E}) & \xrightarrow{\Phi} & \mathfrak{c} \\ \downarrow \pi & & \downarrow \pi & & \downarrow \pi \\ \mathfrak{g} & \xrightarrow{\psi} & \Gamma(TM) & \xrightarrow{\cong} & \text{dif}(M) \end{array}$$

Since the map  $\pi : \mathfrak{c} \longrightarrow \text{dif}(M)$  has a kernel, the algebra of infinitesimal symmetries generated by  $\Psi(\mathfrak{a})$  is, in general, bigger than the corresponding algebra  $\psi(\mathfrak{g})$ . These two will be the same only if  $\Psi(\mathfrak{h})$  acts trivially, i. e.,  $\Psi(\mathfrak{h}) \in \Omega_{cl}^1(M)$ , where  $\mathfrak{h}$  is the kernel of the projection  $\mathfrak{a} \longrightarrow \mathfrak{g}$ . If this is the case, it may still happen that the group of symmetries of  $\mathcal{E}$  generated by  $\Psi(\mathfrak{a})$ ,  $\tilde{G}$ , is bigger than the group of symmetries of  $M$  generated by  $\psi(\mathfrak{g})$ ,  $G$ , as  $\tilde{G}$  may be just a cover of  $G$ .

**Definition 3.6.** An *extended action* is a Courant algebra map  $\Psi : \mathfrak{a} \longrightarrow \Gamma(\mathcal{E})$  which generates the same group of symmetries on  $\mathcal{E}$  and on  $M$ . This means that  $\Psi(\mathfrak{h}) \in \Omega_{cl}^1(M)$  and the infinitesimal actions of  $\Phi \circ \Psi(\mathfrak{a}/\mathfrak{h})$  and  $\psi(\mathfrak{g})$  integrate to an action of the same group.

A practical way to check whether a map of Courant algebras is an extended action is to choose an isotropic splitting for  $\mathcal{E}$  making it

isomorphic to  $(TM \oplus T^*M, \langle \cdot, \cdot \rangle, [\cdot, \cdot]_H)$  and then requiring that this splitting is preserved, i. e.,

$$[\Psi(a), \Gamma(TM)] \subset \Gamma(TM), \quad \text{for all } a \in \mathfrak{a}.$$

Letting  $\Psi(\alpha) = X_\alpha + \xi_\alpha$ , this is equivalent to the condition

$$i_{X_\alpha} H = d\xi_\alpha, \quad \text{for all } \alpha \in \mathfrak{a}. \quad (3.5)$$

If this is the case, then the group action on  $TM \oplus T^*M$  is the one induced by its action on  $TM$  and  $T^*M$  determined by the underlying diffeomorphisms. Reciprocally, if an extended action induces the action of a compact group on  $\mathcal{E}$ , then there is an isotropic splitting preserved by the extended action.

**Proposition 3.7** (Busztyn–Cavalcanti–Gualtieri [11]). *Let the Lie group  $G$  act on the manifold  $M$ , and let  $\mathfrak{a} \xrightarrow{\pi} \mathfrak{g}$  be an exact Courant algebra with a morphism  $\Psi$  to an exact Courant algebroid  $\mathcal{E}$  over  $M$  such that  $\nu(\mathfrak{h}) \subset \Omega_{cl}^1(M)$ .*

*If  $\mathcal{E}$  has a  $\mathfrak{g}$ -invariant splitting, then the  $\mathfrak{g}$ -action on  $\mathcal{E}$  integrates to an action of  $G$ , and hence  $\Psi$  is an extended action of  $G$  on  $\mathcal{E}$ . Conversely, if  $G$  is compact and  $\Psi$  is an extended action, then by averaging splittings one can always find a  $\mathfrak{g}$ -invariant splitting of  $\mathcal{E}$ .*

### 3.1.3 Moment maps

Suppose that we have an extended  $G$ -action on an exact Courant algebroid as in the previous section, so that we have the map  $\nu : \mathfrak{h} \longrightarrow \Omega_{cl}^1(M)$ . Because the action is a Courant algebra morphism, this map is  $\mathfrak{g}$ -equivariant in the sense

$$\nu(g \cdot h) = \mathcal{L}_{\psi(g)} \nu(h).$$

Therefore we are led naturally to the definition of a moment map for this extended action, as an equivariant factorization of  $\mu$  through the smooth functions.

**Definition 3.8.** A *moment map* for an extended  $\mathfrak{g}$ -action on an exact Courant algebroid is a  $\mathfrak{g}$ -equivariant map  $\mu : \mathfrak{h} \longrightarrow C^\infty(M, \mathbb{R})$

satisfying  $d \circ \mu = \nu$ , i.e. such that the following diagram commutes:

$$\begin{array}{ccc}
 & & \mathfrak{h} \\
 & \swarrow \mu & \downarrow \nu \\
 C^\infty(M) & \xrightarrow{d} & \Gamma(T^*M)
 \end{array}$$

Note that  $\mu$  may be alternatively viewed as an equivariant map  $\mu : M \rightarrow \mathfrak{h}^*$ .

As we see next, the usual notions of symplectic and Hamiltonian actions fit into the framework of extended actions of Courant algebras.

**Example 3.9** (Symplectic action). Let  $(M, \omega)$  be a symplectic manifold,  $\mathcal{E}$  be  $(TM \oplus T^*M, \langle \cdot, \cdot \rangle, [\cdot, \cdot])$ , with  $H = 0$ , and  $G$  be a connected Lie group acting on  $M$  by symplectomorphisms, with infinitesimal action  $\psi : \mathfrak{g} \rightarrow \Gamma(TM)$ . Using the adjoint action of  $\mathfrak{g}$  on itself we can regard  $\mathfrak{g}$  as a  $\mathfrak{g}$ -module and using Example 3.3 we have a Courant algebra structure on  $\mathfrak{g} \oplus \mathfrak{g}$ :

$$(g_1, h_1) \bullet (g_2, h_2) = ([g_1, g_2], [g_1, h_2]).$$

If we define  $\Psi : \mathfrak{g} \oplus \mathfrak{g} \rightarrow \Gamma(TM \oplus T^*M)$  by

$$\Psi(g, h) = X_g + \omega(X_h),$$

where  $X_g = \psi(g)$ . Then  $\Psi(g, h)$  satisfies condition (3.5), since in this case  $H = 0$  and  $X_h$  preserves  $\omega$  we have

$$d(\omega(X_h)) = \mathcal{L}_{X_h}\omega - i_{X_h}d\omega = 0 = i_{X_g}H.$$

And finally,  $\Psi$  is a map of Courant algebras, since

$$\begin{aligned}
 \Psi(g_1, h_1) \bullet \Psi(g_2, h_2) &= [X_{g_1}, X_{g_2}] + \mathcal{L}_{X_{g_1}}\omega(X_{h_2}) \\
 &= X_{[g_1, g_2]} + \omega(X_{[g_1, h_2]}) \\
 &= \Psi((g_1, h_1) \bullet (g_2, h_2)).
 \end{aligned}$$

Therefore  $\Psi$  determines an extended action.

The condition for this extended action to have a moment map is that one can find an equivariant map  $\mu : M \longrightarrow \mathfrak{g}^*$  such that

$$\omega(X_h) = d\langle \mu, h \rangle, \quad \forall h \in \mathfrak{g}.$$

This is precisely the moment map condition for symplectic actions, so in this case our definition of moment map is nothing but the usual one.

The analogy with symplectic geometry is very useful not only when dealing with extended actions, but also for the reduction procedure later on. However one particular feature is lost in it: the fact the the group action does not determine the moment map. Next example has overall the same features of the symplectic one, but does not rely on any structure on  $M$ .

**Example 3.10.** Given a Lie algebra  $\mathfrak{g}$ , we can always think of  $\mathfrak{g}$  as a Courant algebra over itself, with the projection given by the identity and the Courant bracket given by the Lie bracket. If  $G$  acts on a manifold  $M$  with infinitesimal action  $\psi : \mathfrak{g} \longrightarrow \Gamma(TM)$  and  $\mathcal{E}$  is an exact Courant algebroid over  $M$ , we can always try and *lift* this action to an extended action on  $\mathcal{E}$ :

$$\begin{array}{ccc} \mathfrak{g} & \xrightarrow{\text{Id}} & \mathfrak{g} \\ \downarrow \tilde{\Psi} & & \downarrow \psi \\ \Gamma(\mathcal{E}) & \xrightarrow{\pi} & \Gamma(TM). \end{array}$$

We call any such an extended action  $\tilde{\Psi}$  a *lifted* or a *trivially extended* action.

If we are also given an equivariant map  $\mu : M \longrightarrow \mathfrak{h}^*$ , where  $\mathfrak{h}$  is a  $\mathfrak{g}$ -module, then we can extend the action  $\tilde{\Psi} : \mathfrak{g} \longrightarrow \Gamma(\mathcal{E})$  to an action of  $\mathfrak{a} = \mathfrak{g} \oplus \mathfrak{h}$ , endowed with the demisemidirect product structure from Example 3.3, by defining

$$\Psi(g, h) = \tilde{\Psi}(g) + d\langle \mu, h \rangle.$$

Similarly to the symplectic case, the equivariance of  $\mu$  implies that this is an extended action.

## 3.2 Reduction of Courant algebroids

In the previous section we saw how a  $G$ -action on a manifold  $M$  could be extended to a Courant algebroid  $\mathcal{E}$ , making it an equivariant  $G$ -bundle in such a way that the Courant structure is preserved by the  $G$ -action. In this section we will see that an extended action determines the reduced Courant algebroid in a more subtle way, as not only does it furnish us the  $G$ -action but also an equivariant subbundle whose quotient is the reduced Courant algebroid. This reduced Courant algebroid is defined over a *reduced manifold* which is the quotient by  $G$  of a submanifold  $P \hookrightarrow M$ , which is also determined by the extended action.

Given an extended action, we have three distributions on  $\mathcal{E}$  associated to it:  $K = \Psi(\mathfrak{a})$ ,  $K^\perp$  and  $K + K^\perp$  and from these we get three distributions on  $M$ :  $\pi(K)$  — the directions of the group action on  $M$  —,  $\Delta_s = \pi(K^\perp)$  and  $\Delta_b = \pi(K + K^\perp)$ . The distribution  $\Delta_s$  can also be described as

$$\Delta_s = \text{Ann}(\Psi(\mathfrak{h})),$$

and hence, since it is the annihilator of a space generated by closed forms, it is locally integrable around the points where  $\Psi(\mathfrak{h})$  has constant rank. Then the leafs of  $\Delta_b$  are just the  $G$ -orbits of the leafs of  $\Delta_s$ . If we pick a leaf  $P$  of  $\Delta_b$  where the group acts freely and properly and over which  $\Psi(\mathfrak{h})$  has constant rank, the reduction procedure can be described using the following facts:

1. The Courant bracket on  $\mathcal{E}$  induces a bracket on  $\Gamma(K^\perp|_P)^G$  which is well defined modulo  $\Gamma(K \cap K^\perp|_P)^G$ ,
2. Using the bracket above in  $\Gamma(K^\perp|_P)^G$ , we have that  $\Gamma(K \cap K^\perp|_P)^G$  is an ideal, so this bracket induces a well defined bracket on

$$\mathcal{E}_{red} = \frac{K^\perp}{K \cap K^\perp} \Big/ G \quad \text{over} \quad M_{red} = P/G.$$

This means that given two sections  $e_1, e_2 \in \Gamma(\mathcal{E}_{red})$ , their bracket is defined by choosing  $G$ -invariant lifts  $\tilde{e}_1, \tilde{e}_2 \in \Gamma(\mathcal{E})|_P$ , then extending

these lifts *arbitrarily* to sections of  $\mathbf{e}_1, \mathbf{e}_2 \in \Gamma(\mathcal{E})$  and then letting

$$\llbracket e_1, e_2 \rrbracket := \llbracket \mathbf{e}_1, \mathbf{e}_2 \rrbracket|_P + K \cap K^\perp \subset \frac{K^\perp}{K \cap K^\perp}.$$

According to (1) and (2) above, this bracket is independent of the choices made. It is clear that  $\mathcal{E}_{red}$  comes equipped with a projection  $\mathcal{E}_{red} \rightarrow TM_{red}$ . Further,  $K \cap K^\perp$  is the null space of the pairing restricted to  $K^\perp$  and hence  $E_{red}$  has a pairing. Together, the bracket, the projection and the pairing give  $\mathcal{E}_{red}$  the structure of a Courant algebroid over  $M_{red}$ .

We call  $M_{red}$  a *reduced manifold* and  $\mathcal{E}_{red}$  the *reduced Courant algebroid* over it. The argument above is the main ingredient in the proof of the following theorem.

**Theorem 3.11** (Burzstyn–Cavalcanti–Gualtieri [11]). *Given an extended action  $\Psi$ , let  $P$  be a smooth leaf of the distribution  $\Delta_b$  where  $\Psi(\mathfrak{h})$  has constant rank. Then the vector bundle  $\mathcal{E}_{red}$  defined above is a Courant algebroid over  $M_{red} = P/G$ .  $\mathcal{E}_{red}$  is exact if and only if*

$$\pi(K) \cap \pi(K^\perp) = \pi(K \cap K^\perp) \text{ over } P.$$

*In particular  $\mathcal{E}_{red}$  is exact if  $K$  is isotropic over  $P$ .*

In all the cases we will treat where  $\mathcal{E}_{red}$  is exact,  $K$  will be isotropic over  $P$ . As we will see, in the presence of a moment map, this is an analogue of requiring that  $M_{red}$  is the inverse image of zero.

### 3.2.1 Examples

In this section we will provide some examples of Courant algebroid reduction. Since Courant algebroids are often given together with a splitting, we describe the behaviour of splittings under reduction. This is then related to the way in which the Ševera class  $[H]$  of an exact Courant algebroid is transported to the reduced space.

**Example 3.12.** Assume  $G$  acts freely and properly on  $M$ , with infinitesimal action  $\psi : \mathfrak{g} \rightarrow \Gamma(TM)$  and consider the split Courant algebroid  $(TM \oplus T^*M, \langle \cdot, \cdot \rangle, \llbracket \cdot, \cdot \rrbracket_H)$ , where  $H$  is a basic 3-form.

Using the inclusion  $TM \rightarrow TM \oplus T^*M$  we obtain an action of the Courant algebra  $\mathfrak{a} = \mathfrak{g}$  which preserves the splitting (condition

(3.5) holds as  $H$  is basic). In this case  $K^\perp = TM \oplus \text{Ann}(\psi(\mathfrak{g}))$  and hence  $\Delta_b = \pi(K^\perp + K) = TM$  has only one leaf:  $M$ .

Therefore the only possible reduced manifold in this case is  $M^{red} = M/G$  and the reduced algebroid is

$$\mathcal{E}^{red} = \frac{K^\perp/G}{K/G} = TM/\psi(\mathfrak{g}) \oplus \text{Ann}(\psi(\mathfrak{g})) \cong TM^{red} \oplus T^*M^{red}.$$

Since  $H$  is a basic form, it is a well defined form on  $M^{red}$  and one can easily check that  $H$  is the curvature of the reduced algebroid for this splitting.

**Example 3.13.** Even a trivial group action may be extended by 1-forms. Consider the extended action  $\rho : \mathbb{R} \rightarrow \Gamma(\mathcal{E})$  on an exact Courant algebroid  $\mathcal{E}$  over  $M$  given by  $\rho(1) = \xi$  for some closed 1-form  $\xi$ . Then  $K = \langle \xi \rangle$  and  $K^\perp = \{v \in \mathcal{E} : \pi(v) \in \text{Ann}(\xi)\}$  which induces the distribution  $\Delta_b = \text{Ann}(\xi) \subset TM$ , which is integrable wherever  $\xi$  is nonzero. Since the group action is trivial, a reduced manifold is simply a choice of integral submanifold  $\iota : P \hookrightarrow M$  for  $\xi$  and the reduced Courant algebroid is just the restricted Courant algebroid as introduced in Section 1.5. The characteristic class of  $\mathcal{E}^{red}$  in this case is the pullback to  $P$  of the class of  $\mathcal{E}$ .

In the preceding examples, the reduced Courant algebroid inherited a natural splitting; this is not always the case. The next example demonstrates this as well as the phenomenon by which a trivial twisting  $[H] = 0$  may give rise to a reduced Courant algebroid with nontrivial curvature.

**Example 3.14.** Assume that  $S^1$  acts freely and properly on  $M$  with infinitesimal action  $\psi : \mathfrak{s}^1 \rightarrow \Gamma(TM)$ ,  $\psi(1) = \partial_\theta$ , and let  $\Psi : \mathfrak{s}^1 \rightarrow \Gamma(\mathcal{E})$  be a trivial extension of this action (i.e.,  $\mathfrak{h} = \{0\}$ ) such that  $\Psi(\mathfrak{s}^1) = K$  is isotropic. In these conditions  $\Delta_s = \text{Ann}(\mathfrak{h}) = TM$ , so the only leaf of  $\Delta_s$  is  $M$  itself and  $M_{red} = M/S^1$ .

By Proposition 3.7, we may choose an invariant splitting so that  $\mathcal{E} = (TM \oplus T^*M, \langle \cdot, \cdot \rangle, [\cdot, \cdot]_H)$ , with  $\Psi(1) = \partial_\theta + \xi$  and

$$i_{\partial_\theta} H = d\xi. \tag{3.6}$$

The form  $\xi$  is basic as isotropy implies that  $\xi(\partial_\theta) = 0$  and condition (3.6) tells us it is invariant:

$$\mathcal{L}_{\partial_\theta}\xi = d\xi(\partial_\theta) + i_{\partial_\theta}d\xi = i_{\partial_\theta}i_{\partial_\theta}H = 0.$$

Equation (3.6) also implies that  $H$  is invariant under the circle action.

Now we show that if we choose a connection  $\theta$  for the circle bundle  $M$ , then we can reduce this example to Example 3.12 via a nonclosed  $B$ -field transform. Indeed, once we have chosen  $\theta$ , we have

$$H = d\xi \wedge \theta + h,$$

where  $\xi$  and  $h$  are basic forms.

We use the connection  $\theta$  to change the splitting of  $TM \oplus T^*M$  by the nonclosed 2-form  $B = \theta \wedge \xi$ . In this new splitting the action is given by

$$\partial_\theta + \xi - i_{\partial_\theta}\langle\theta, \xi\rangle = \partial_\theta.$$

And the curvature of the new splitting is given by

$$H^{red} = H + dB = \theta \wedge d\xi + h + d(\theta \wedge \xi) = \langle d\theta, \xi \rangle + h. \quad (3.7)$$

Observe that  $H^{red}$  is basic, and therefore, according to Example 3.12, is the curvature of the reduced algebroid over  $M/G$ .

Observe that the splitting of the reduced Courant algebroid over  $M_{red}$  obtained above is not determined by the original splitting of  $\mathcal{E}$  alone, but also the choice of connection  $\theta$ . Also, if  $H = 0$ , the curvature of the reduced algebroid is given by  $F \wedge \xi$ , which may be a nontrivial cohomology class on  $M_{red}$ , therefore showing that even if  $\mathcal{E}$  has trivial characteristic class  $\mathcal{E}_{red}$  can have nonvanishing characteristic class.

**Exercise 3.15.** Assume that  $\mathcal{E}$  is equipped with a  $G$ -invariant splitting  $\nabla$  and *the action*  $\Psi$  is *split*, in the sense that there is a splitting  $s$  for  $\pi : \mathfrak{a} \rightarrow \mathfrak{g}$  making the diagram commutative:

$$\begin{array}{ccc} \mathfrak{a} & \xleftarrow{s} & \mathfrak{g} \\ \downarrow \Psi & & \downarrow \psi \\ \Gamma(\mathcal{E}) & \xleftarrow{\nabla} & \Gamma(TM) \end{array} \quad (3.8)$$

Show that in this case  $\mathcal{E}_{red}$  is exact and has a natural splitting.



**Example 3.16** (Symplectic reduction I). Let  $(M, \omega)$  be a symplectic manifold and consider the extended  $G$ -action  $\Psi : \mathfrak{g} \oplus \mathfrak{g} \longrightarrow \Gamma(TM \oplus T^*M)$  with curvature  $H = 0$  defined in Example 3.9.

Let  $\psi : \mathfrak{g} \longrightarrow \Gamma(TM)$  be the infinitesimal action and  $\psi(\mathfrak{g})^\omega$  denote the symplectic orthogonal of the image distribution  $\psi(\mathfrak{g})$ . Then the extended action has image

$$K = \psi(\mathfrak{g}) \oplus \omega(\psi(\mathfrak{g})),$$

so that the orthogonal complement is

$$K^\perp = \psi(\mathfrak{g})^\omega \oplus \text{Ann}(\psi(\mathfrak{g})).$$

Then the distributions  $\Delta_b$  and  $\Delta_s$  on  $M$  are

$$\begin{aligned} \Delta_b &= \psi(\mathfrak{g})^\omega + \psi(\mathfrak{g}), \\ \Delta_s &= \psi(\mathfrak{g})^\omega. \end{aligned}$$

If the action is Hamiltonian, with moment map  $\mu : M \longrightarrow \mathfrak{g}^*$ , then  $\Delta_s$  is the tangent distribution to the level sets  $\mu^{-1}(\lambda)$  while  $\Delta_b$  is the tangent distribution to the sets  $\mu^{-1}(\mathcal{O}_\lambda)$ , for  $\mathcal{O}_\lambda$  a coadjoint orbit containing  $\lambda$ . Therefore we see that the reduced Courant algebroid is simply  $TM_{red} \oplus T^*M_{red}$  with  $H = 0$ , for the usual symplectic reduced space  $M_{red} = \mu^{-1}(\mathcal{O}_\lambda)/G = \mu^{-1}(\lambda)/G_\lambda$ .

We finish with an example which combines features of Examples 3.10, 3.14 and 3.16

**Example 3.17.** Let  $\tilde{\Psi} : \mathfrak{g} \longrightarrow \Gamma(\mathcal{E})$  be an isotropic trivially extended action

$$\begin{array}{ccc} \mathfrak{g} & \xrightarrow{\text{Id}} & \mathfrak{g} \\ \downarrow \tilde{\Psi} & & \downarrow \psi \\ \Gamma(\mathcal{E}) & \xrightarrow{\pi} & \Gamma(TM) \end{array}$$

and let  $\mu : M \longrightarrow \mathfrak{h}^*$  be an equivariant map, where  $\mathfrak{h}$  is a  $\mathfrak{g}$ -module. Then, according to Example 3.10 we can extend the action  $\tilde{\Psi} : \mathfrak{g} \longrightarrow \Gamma(\mathcal{E})$  to an action of  $\mathfrak{a} = \mathfrak{g} \oplus \mathfrak{h}$ , endowed with the hemisemidirect product structure from Example 3.3, by defining

$$\Psi(g, h) = \tilde{\Psi}(g) + d\langle \mu, h \rangle.$$

The reduced manifolds for the action  $\Psi$  correspond to  $\mu^{-1}(\mathcal{O}_\lambda)/G$ , where  $\mathcal{O}_\lambda$  is the  $G$ -orbit of  $\lambda \in \mathfrak{h}^*$ . In this setting,  $K$  is only isotropic over  $P = \mu^{-1}(\lambda)$  with  $\lambda$  a central element in  $\mathfrak{h}^*$ , so the requirement that  $K$  is isotropic is an analogue of the condition  $P = \mu^{-1}(0)$ .

In order to describe the reduced algebroid over  $\mu^{-1}(0)$  we observe that the reduction can be described in two steps. The first is just restriction to the level set  $P = \mu^{-1}(0)$ . Then the extended action on  $M$  gives rise to an isotropic trivially extended action on  $P$  and the second step is to perform the reduction by this action.

If we have an invariant splitting  $\mathcal{E} = TM \oplus T^*M$  with the  $H$ -bracket, i. e., equation (3.5) holds for this splitting, then the first step gives as intermediate Courant algebroid  $TP \oplus T^*P$  with the  $\iota^*H$ -bracket, where  $\iota : P \hookrightarrow M$  is the inclusion map and we have an extended action  $\mathfrak{g} \longrightarrow \Gamma(TP \oplus T^*P)$

$$g \mapsto X_g + \xi_g.$$

satisfying:

$$i_{X_g}H|_P = d\xi_g. \quad (3.5)$$

In particular, projecting onto  $T^*P$ , we obtain  $\xi \in \Omega^1(P; \mathfrak{g}^*)$ . According to our general assumptions,  $G$  acts freely and properly on  $P$ , making it a principal  $G$ -bundle, so we can repeat the argument from Example 3.14: Let  $\theta \in \Omega^1(P; \mathfrak{g})$  be a connection for this bundle and consider the change of splitting  $TP \oplus T^*P$  determined by the 2-form  $B = \langle \theta, \xi \rangle$ , where  $\langle \cdot, \cdot \rangle$  denotes the pairing between  $\mathfrak{g}$  and  $\mathfrak{g}^*$  together with the wedge product of forms. In this new splitting the action is given by

$$g \mapsto X_g + \xi_g - i_{X_g}\langle \theta, \xi \rangle = X_g.$$

Further, according to (3.5) we can write the invariant form  $\iota^*H$  as

$$\iota^*H = \langle \theta, d\xi \rangle + h,$$

with  $h \in \Omega^3(P/G)$  a basic form. Hence, the curvature of the new splitting is given by

$$H^{red} = \iota^*H + dB = \langle \theta, d\xi \rangle + h + \langle d\theta, \xi \rangle - \langle \theta, d\xi \rangle = \langle d\theta, \xi \rangle + h.$$

Observe that  $H^{red}$  is basic, and therefore, according to Example 3.12, is the curvature of the reduced algebroid over  $P/G$ .

### 3.3 Reduction of Dirac structures

In this section we study how to transport Dirac structures invariant under an extended action from  $\mathcal{E}$  to  $\mathcal{E}_{red}$ .

The basic observation which allows us to reduce Dirac structures is a simple piece of linear algebra: Given an isotropic subspace  $K \subset \mathcal{D}V$ , it furnishes a way to transport linear Dirac structures on  $\mathcal{D}V$  to linear Dirac structures on  $K^\perp/K$ . If  $D \subset \mathcal{D}V$  is a Dirac structure, we define

$$D_{red} = \frac{D \cap K^\perp + K}{K}.$$

And  $D_{red}$  is a Dirac structure on  $K^\perp/K$  since

$$D_{red}^\perp = \frac{(D \cap K^\perp + K)^\perp}{K} = \frac{(D + K) \cap K^\perp}{K} = \frac{D \cap K^\perp + K}{K} = D.$$

Observe that this idea was used before when studying generalized complex submanifolds.

Now, we move on to the reduction procedure and let  $\Psi : \mathfrak{a} \rightarrow \Gamma(\mathcal{E})$  be an extended action for which the reduced Courant algebroid over a reduced manifold  $M_{red}$  is exact. If a Dirac structure  $D$  is preserved by  $\Psi$ , i.e.,  $\Psi(\mathfrak{a}) \bullet \Gamma(D) \subset \Gamma(D)$ , then we have a natural candidate for a Dirac structure on the reduced algebroid:

$$D_{red} = \frac{(D \cap K^\perp + K)^G}{K^G} \Big|_{M_{red}} \subset \mathcal{E}_{red}. \quad (3.9)$$

**Theorem 3.18** (Bursztyn–Cavalcanti–Gualtieri [11]). *Let  $\rho : \mathfrak{a} \rightarrow \Gamma(\mathcal{E})$  be an extended action preserving a Dirac structure  $D \subset \mathcal{E}$  such that  $K$  is isotropic over  $P$ , a leaf of  $\Delta_b$ . If  $(D \cap K)|_P$  is a smooth bundle, then*

$$D_{red} = \frac{(D \cap K^\perp + K)^G}{K^G} \Big|_{M_{red}} \subset \mathcal{E}_{red}. \quad (3.10)$$

defines a Dirac structure on  $\mathcal{E}_{red}$ .

*Proof.* The distribution  $D_{red}$  is certainly maximal isotropic and, since  $D \cap K|_P$  is a smooth bundle,  $D_{red}$  is just the smooth quotient of two

bundles and hence is smooth. Given two sections  $e_1, e_2 \in \Gamma(D_{red})$ , let  $\tilde{e}_1, \tilde{e}_2 \in \Gamma((D \cap K^\perp + K)|_P)$  be  $G$ -invariant representatives of them in  $K^\perp G$ . Then we can write

$$\tilde{e}_i = \tilde{e}_i^D + \tilde{e}_i^K,$$

so that  $\tilde{e}_i^D \in \Gamma(D \cap K^\perp|_P)$  and  $\tilde{e}_i^K \in \Gamma(K|_P)$  are smooth sections.

Now extend  $\tilde{e}_i^D$  to invariant sections of  $D \cap K^\perp$  and  $\tilde{e}_i^K$  to invariant sections of  $K$ , so that

$$[[\tilde{e}_1, \tilde{e}_2]] = [[e_1^D, e_2^D]] + [[e_1^D, e_2^K]] + [[e_1^K, e_2^D]] + [[e_1^K, e_2^K]].$$

The first term above lies in  $D \cap K^\perp$ , since both  $D$  and  $\Gamma(K^\perp)^G$  are closed under the bracket and  $e_i^D \in \Gamma(D \cap K^\perp)^G$ . The remaining terms lie in  $\Gamma(K)^G$ , since this is an ideal of  $\Gamma(K^\perp)^G$ . Therefore

$$[[e_1, e_2]] = [[\tilde{e}_1, \tilde{e}_2]]|_P + K \subset \Gamma(D \cap K^\perp + K)^G,$$

showing that  $D_{red}$  is closed under the bracket in  $\mathcal{E}_{red}$ .  $\square$

The reduction of Dirac structures works in the same way for *complex* Dirac structures, provided one replaces  $K$  by its complexification.

### 3.4 Reduction of generalized complex structures

As we know, a generalized complex structure is a complex Dirac structure  $L \subset \mathcal{E}_\mathbb{C}$  satisfying  $L \cap \overline{L} = \{0\}$ . So, if an extended action  $\Psi$  preserves a generalized complex structure  $\mathcal{J}$  with  $i$ -eigenspace  $L$  and we pick a leaf  $P$  of the distribution  $\Delta_b$  over which  $K$  is isotropic, we can try and reduce  $L$ , as a Dirac structure:

$$L_{red} = \left. \frac{(L \cap K_\mathbb{C}^\perp + K_\mathbb{C})^G}{K_\mathbb{C}^G} \right|_{M_{red}} \quad (3.11)$$

This reduced Dirac structure  $L_{red}$  is not necessarily a generalized complex structure as it may not satisfy  $L_{red} \cap \overline{L_{red}} = \{0\}$ . Whenever it does it determines a generalized complex structure.

The condition  $L_{red} \cap \overline{L_{red}} = \{0\}$  is a simple linear algebraic condition which can be rephrased in the following way (compare with the condition for a submanifold to be a generalized complex submanifold).

**Lemma 3.19.** *The distribution  $L_{red}$  satisfies  $L_{red} \cap \overline{L_{red}} = \{0\}$  if and only if*

$$\mathcal{J}K \cap K^\perp \subset K \text{ over } P. \quad (3.12)$$

*Proof.* It is clear from (3.11) that  $L_{red} \cap \overline{L_{red}} = \{0\}$  over the reduced manifold if and only if

$$(L \cap K_{\mathbb{C}}^\perp + K_{\mathbb{C}}) \cap (\overline{L} \cap K_{\mathbb{C}}^\perp + K_{\mathbb{C}}) \subset K_{\mathbb{C}} \text{ over } P. \quad (3.13)$$

Hence, we must prove that conditions (3.12) and (3.13) are equivalent.

We first prove that (3.12) implies (3.13). Let  $v \in (L \cap K_{\mathbb{C}}^\perp + K_{\mathbb{C}}) \cap (\overline{L} \cap K_{\mathbb{C}}^\perp + K_{\mathbb{C}})$  over a given point. Without loss of generality we can assume that  $v$  is real. Since  $v \in L \cap K_{\mathbb{C}}^\perp + K_{\mathbb{C}}$ , we can find  $v_L \in L \cap K_{\mathbb{C}}^\perp$  and  $v_K \in K_{\mathbb{C}}$  such that  $v = v_L + v_K$ . Taking conjugates, we get that  $v = \overline{v_L} + \overline{v_K}$ , hence  $v_L - \overline{v_L} = \overline{v_K} - v_K$ . Applying  $-i\mathcal{J}$ , we obtain

$$v_L + \overline{v_L} = -i\mathcal{J}(\overline{v_K} - v_K).$$

The left hand side lies in  $K^\perp$  while the right hand side lies in  $\mathcal{J}K$ . It follows from (3.12) that  $v_L + \overline{v_L} \in K$ , hence  $v = \frac{1}{2}(v_L + \overline{v_L} + v_K + \overline{v_K}) \in K$ , as desired.

Conversely, if (3.12) does not hold, i.e., there is  $v \in \mathcal{J}K \cap K^\perp$  with  $v \notin K$ , then  $v - i\mathcal{J}v \in L \cap K_{\mathbb{C}}^\perp$  and  $v + i\mathcal{J}v \in \overline{L} \cap K_{\mathbb{C}}^\perp$ . Since  $v \in \mathcal{J}K$  and  $\mathcal{J}v \in K$ , it follows that  $v \in L \cap K_{\mathbb{C}}^\perp + K_{\mathbb{C}}$  and  $v \in \overline{L} \cap K_{\mathbb{C}}^\perp + K_{\mathbb{C}}$ , showing that  $(L \cap K_{\mathbb{C}}^\perp + K_{\mathbb{C}}) \cap (\overline{L} \cap K_{\mathbb{C}}^\perp + K_{\mathbb{C}}) \not\subset K_{\mathbb{C}}$ . This concludes the proof.  $\square$

So, this lemma tells us precisely when a generalized complex structure can be reduced. However (3.12) may be hard to check in real examples, so we settle with more meaningful conditions in the following theorems.

**Theorem 3.20** (Bursztyn–Cavalcanti–Gualtieri [11]). *Let  $\Psi$  be an extended  $G$ -action on the exact Courant algebroid  $\mathcal{E}$ . Let  $P$  be a leaf of the distribution  $\Delta_b$  over which  $K$  is isotropic and where  $G$  acts freely and properly. If the action preserves a generalized complex structure  $\mathcal{J}$  on  $\mathcal{E}$  and  $\mathcal{J}K = K$  over  $P$  then  $\mathcal{J}$  reduces to  $\mathcal{E}_{red}$ .*

*Proof.* Since  $K$  is isotropic over  $P$ , the reduced algebroid is exact. Further, the condition  $\mathcal{J}K = K$  implies that  $L \cap K_{\mathbb{C}}$  is smooth, as it is just the  $i$ -eigenspace of  $\mathcal{J}|_K$ . So, according to Theorem 3.18,  $L$  reduces as a Dirac structure. Finally,

$$\mathcal{J}K \cap K^{\perp} = K \cap K^{\perp} = K,$$

and hence Lemma 3.19 implies that the reduced Dirac structure is a generalized complex structure.  $\square$

*Remark:* The theorem still holds even if  $K$  is not isotropic over  $P$ , but as long as the reduced algebroid is exact [11]. This is the analogue of saying that one can do symplectic reduction for any value of the moment map and not only the inverse image of a central element.

**Corollary 3.21.** *If the hypotheses of the previous theorem hold and the extended action has a moment map  $\mu : M \rightarrow \mathfrak{h}^*$ , then the reduced Courant algebroid over  $\mu^{-1}(O_{\lambda})/G$  has a reduced generalized complex structure.*

It is easy to check that the reduced generalized complex structure  $\mathcal{J}^{red}$  constructed in Theorem 3.20 is characterized by the following commutative diagram:

$$\begin{array}{ccc} K^{\perp} & \xrightarrow{\mathcal{J}} & K^{\perp} \\ \downarrow & & \downarrow \\ \frac{K^{\perp}}{K \cap K^{\perp}} & \xrightarrow{\mathcal{J}^{red}} & \frac{K^{\perp}}{K \cap K^{\perp}} \end{array} \quad (3.14)$$

Theorem 3.20 uses the compatibility condition  $\mathcal{J}K = K$  for the reduction of  $\mathcal{J}$ . We now observe that the reduction procedure also works in an extreme opposite situation.

**Theorem 3.22** (Bursztyn–Cavalcanti–Gualtieri [11]). *Consider an extended  $G$ -action  $\Psi$  on an exact Courant algebroid  $\mathcal{E}$ . Let  $P$  be a leaf of the distribution  $\Delta_b$  where  $G$  acts freely and properly. If  $K$  is isotropic over  $P$  and  $\langle \cdot, \cdot \rangle : K \times \mathcal{J}K \rightarrow \mathbb{R}$  is nondegenerate then  $\mathcal{J}$  reduces.*

*Proof.* As  $K$  is isotropic over  $P$ , the reduced Courant algebroid is exact. The nondegeneracy assumption implies that  $\mathcal{J}K \cap K^\perp = \{0\}$  and in particular  $\mathcal{J}K \cap K = \{0\}$ . The latter implies that  $L \cap K_{\mathbb{C}} = \{0\}$ , which is a smooth bundle, so according to Theorem 3.18,  $L$  reduces as a Dirac structure, while the former implies that Lemma 3.19 holds and therefore the reduced Dirac structure is a generalized complex structure.  $\square$

### 3.4.1 Symplectic structures

We now present two examples of reduction obtained from a symplectic manifold  $(M, \omega)$ : First, we show that ordinary symplectic reduction is a particular case of our construction; the second example illustrates how one can obtain a type 1 generalized complex structure as the reduction of an ordinary symplectic structure. In both examples, the initial Courant algebroid is just  $TM \oplus T^*M$  with  $H = 0$ .

**Example 3.23** (Symplectic reduction II). Let  $(M, \omega)$  be a symplectic manifold, and let  $\mathcal{J}_\omega$  be the generalized complex structure associated with  $\omega$ . Following Example 3.9 and keeping the same notation, consider a symplectic  $G$ -action on  $M$ , regarded as an extended action. It is clear that  $\mathcal{J}_\omega K = K$ , so we are in the situation of Theorem 3.20 and issuing remark.

Following Example 3.16, let  $S$  be a leaf of the distribution  $\Delta_s = \psi(\mathfrak{g})^\omega$ . Since  $K$  splits as  $K_T \oplus K_{T^*}$ , the reduction procedure of Theorem 3.18 in this case amounts to the usual pull-back of  $\omega$  to  $S$ , followed by a Dirac push-forward to  $S/G_s = M_{red}$ . If the symplectic action admits a moment map  $\mu : M \rightarrow \mathfrak{g}^*$ , then the leaves of  $\Delta_s$  are level sets  $\mu^{-1}(\lambda)$ , and Theorem 3.20 simply reproduces the usual Marsden-Weinstein quotient  $\mu^{-1}(\lambda)/G_\lambda$ . The condition that  $K$  is isotropic over  $P$  in this case corresponds to taking a central element  $\lambda$ .

Next, we show that by allowing the projection  $\pi : K \rightarrow TM$  to be injective, one can reduce a symplectic structure (type 0) to a generalized complex structure with nonzero type.

**Example 3.24.** Assume that  $X$  and  $Y$  are linearly independent symplectic vector fields generating a  $T^2$ -action on  $M$ . Assume further

that  $\omega(X, Y) = 0$  and consider the extended  $T^2$ -action on  $T \oplus T^*$  defined by

$$\Psi(\alpha_1) = X + \omega(Y); \quad \Psi(\alpha_2) = -Y + \omega(X),$$

where  $\{\alpha_1, \alpha_2\}$  is the standard basis of  $\mathfrak{t}^2 = \mathbb{R}^2$ . It follows from  $\omega(X, Y) = 0$  and the fact that the vector fields  $X$  and  $Y$  are symplectic that this is an extended action with isotropic  $K$ .

Since  $\mathcal{J}_\omega K = K$ , Theorem 3.20 implies that the quotient  $M/T^2$  has an induced generalized complex structure. Note that

$$L \cap K_{\mathbb{C}}^\perp = \{Z - i\omega(Z) : Z \in \text{Ann}(\omega(X) \wedge \omega(Y))\},$$

and it is simple to check that  $X - i\omega(X) \in L \cap K_{\mathbb{C}}^\perp$  represents a nonzero element in  $L_{red} = ((L \cap K_{\mathbb{C}}^\perp + K_{\mathbb{C}})/K_{\mathbb{C}})/G$  which lies in the kernel of the projection  $L_{red} \rightarrow T(M/T^2)$ . As a result, this reduced generalized complex structure has type 1.

One can find concrete examples illustrating this construction by considering symplectic manifolds which are  $T^2$ -principal bundles with lagrangian fibres, such as  $T^2 \times T^2$ , or the Kodaira–Thurston manifold. In these cases, the reduced generalized complex structure determines a complex structure on the base 2-torus.

### 3.4.2 Complex structures

In this section we show how a complex manifold  $(M, I)$  may have different types of generalized complex reductions.

**Example 3.25** (Holomorphic quotient). Let  $G$  be a complex Lie group acting holomorphically on  $(M, I)$ , so that the induced infinitesimal map  $\Psi : \mathfrak{g} \rightarrow \Gamma(TM)$  is a holomorphic map. Since  $K = \Psi(\mathfrak{g}) \subset TM$ , it is clear that  $K$  is isotropic and the reduced Courant algebroid is exact. Furthermore, as  $\Psi$  is holomorphic, it follows that  $\mathcal{J}_I K = K$ . By Theorem 3.20, the complex structure descends to a generalized complex structure in the reduced manifold  $M/G$ . The reduced generalized complex structure is nothing but the quotient complex structure obtained from holomorphic quotient.

**Exercise 3.26.** Let  $(M, I)$  be a complex manifold,  $\Psi : \mathfrak{a} \rightarrow \Gamma(TM \oplus T^*M)$  be an extended action and  $K = \Psi(\mathfrak{a})$ . If  $\mathcal{J}_I$  is the general-



ized complex structure induced by  $I$ , show that if  $\mathcal{J}_I K = K$ , then reduction of  $\mathcal{J}_I$  is of complex type.

**Example 3.27.** Consider  $\mathbb{C}^2$  equipped with its standard holomorphic coordinates  $(z_1 = x_1 + iy_1, z_2 = x_2 + iy_2)$ , and let  $\Psi$  be the extended  $\mathbb{R}^2$ -action on  $\mathbb{C}^2$  defined by

$$\Psi(\alpha_1) = \partial_{x_1} + dx_2, \quad \Psi(\alpha_2) = \partial_{y_2} + dy_1,$$

where  $\{\alpha_1, \alpha_2\}$  is the standard basis for  $\mathbb{R}^2$ . Note that  $K = \Psi(\mathbb{R}^2)$  is isotropic, so the reduced Courant algebroid over  $\mathbb{C}/\mathbb{R}^2$  is exact. Since the natural pairing between  $K$  and  $\mathcal{J}_I K$  is nondegenerate, Proposition 3.22 implies that one can reduce  $\mathcal{J}_I$  by this extended action. In this example, one computes

$$K_{\mathbb{C}}^{\perp} \cap L = \text{span}\{\partial_{x_1} - i\partial_{x_2} - dy_1 + idx_1, \partial_{y_1} - i\partial_{y_2} - dy_2 + idx_2\}$$

and  $K_{\mathbb{C}}^{\perp} \cap L \cap K_{\mathbb{C}} = \{0\}$ . As a result,  $L_{red} \cong K_{\mathbb{C}}^{\perp} \cap L$ . So  $\pi : L_{red} \rightarrow \mathbb{C}^2/\mathbb{R}^2$  is an injection, and  $\mathcal{J}^{red}$  has zero type, i.e., it is of symplectic type.

# Chapter 4

## T-duality with NS-flux and generalized complex structures

T-duality in physics is a symmetry which relates IIA and IIB string theory and T-duality transformations act on spaces in which at least one direction has the topology of a circle. In this chapter, we consider a mathematical version of T-duality introduced by Bouwknegt, Evslin and Mathai for principal circle bundles with nonzero twisting 3-form  $H$  [7, 8].

From the point of view adopted in these notes, the relation between two T-dual spaces can be best described using the language of Courant algebroids. Two T-dual spaces are principal circle bundles  $E$  and  $\tilde{E}$  over a common base  $M$  and hence the space of invariant sections of  $(TE \oplus T^*E, \langle \cdot, \cdot \rangle, [\cdot, \cdot]_H)$  and  $(T\tilde{E} \oplus T^*\tilde{E}, \langle \cdot, \cdot \rangle, [\cdot, \cdot]_{\tilde{H}})$  can both be identified with nonexact Courant algebroids over  $M$ . With that said, the T-duality condition is nothing but requiring that these Courant algebroids are isomorphic:

$$\begin{array}{ccc} (TE \oplus T^*E)^{S^1}, \langle \cdot, \cdot \rangle, [\cdot, \cdot]_H & \xrightarrow{\cong} & (T\tilde{E} \oplus T^*\tilde{E})^{S^1}, \langle \cdot, \cdot \rangle, [\cdot, \cdot]_{\tilde{H}} \\ & \searrow \pi & \swarrow \tilde{\pi} \\ & TM & \end{array}$$

Therefore any invariant structure on  $TE \oplus T^*E$  can be transported to an invariant structure on  $T\tilde{E} \oplus T^*\tilde{E}$ . This is particularly interesting since  $E$  and  $\tilde{E}$  have different topologies, in general. Further, when using the map above to transport generalized complex structures, the type changes by  $\pm 1$ . This means that even if  $E$  is endowed with a symplectic or complex structure, the corresponding structure on  $\tilde{E}$  will not be either complex or symplectic, but just generalized complex.

Another structure which can be transported by the isomorphism above is a generalized metric invariant under the circle action. Since a generalized metric can be described in terms of a metric  $g$  on  $E$  and a 2-form  $b$ , studying the way the generalized metric transforms is equivalent to studying the transformations rules for  $g$  and  $b$ . As we will see, these rules are nothing but the Buscher rules [13, 14], which are obtained in a geometrical way, using this point of view.

A final interesting point is that given two principal circle bundles  $E$  and  $\tilde{E}$  over  $M$ , we can always form the fiber product, or *correspondence space*,  $E \times_M \tilde{E}$ , which we can endow with, say, the 3-form  $H$  from  $E$ . The condition that  $E$  and  $\tilde{E}$  are T-dual can then be stated by saying that  $(TE \oplus T^*E, \langle \cdot, \cdot \rangle, \llbracket \cdot, \cdot \rrbracket_H)$  and  $(T\tilde{E} \oplus T^*\tilde{E}, \langle \cdot, \cdot \rangle, \llbracket \cdot, \cdot \rrbracket_{\tilde{H}})$  are different reductions of the Courant algebroid over the correspondence space with curvature  $H$ :

$$\begin{array}{ccc}
 & (E \times_M \tilde{E}, p^*H) & \\
 \swarrow / \partial_{\tilde{\theta}} & & \searrow / \frac{\partial}{\partial \theta} - \tilde{\theta} \\
 (E, H) & & (\tilde{E}, \tilde{H})
 \end{array}$$

This chapter is heavily based on a collaborative work with Gualtieri [18] and organized in the following way. In the first section we introduce T-duality for principal circle bundles as presented in [7] and prove the main result in that paper stating that T-dual manifolds have isomorphic twisted cohomologies. In Section 4.2 we prove that T-duality can be expressed as an isomorphism of Courant algebroids and hence Dirac and generalized complex structures can be transported via T-duality as well as a generalized metric. In the last

section we show that T-duality can be seen in the light of reduction of Courant algebroids.

## 4.1 T-duality with NS-flux

In this section we review the definition of T-duality for principal circle bundles as expressed by Bouwknegt, Evslin and Mathai [7] and some of their results regarding T-dual spaces.

Given a principal circle bundle  $E \xrightarrow{\pi} M$ , with an invariant closed integral 3-form  $H \in \Omega^3(E)$  and a connection  $\theta$ , we can always write  $H = \tilde{F} \wedge \theta + h$ , where  $\tilde{F}$  and  $h$  are basic forms. We denote by  $F = d\theta$  the curvature of  $\theta$ . Bouwknegt *et al* define the *T-dual space* to be another principal circle bundle  $\tilde{E}$  over  $M$  with a connection  $\tilde{\theta}$  whose curvature is the pushforward of  $H$  to  $M$ ,  $d\tilde{\theta} = \pi_! H = \tilde{F}$ , (and this determines  $\tilde{E}$ ) with associated 3-form  $\tilde{H} = F \wedge \tilde{\theta} + h$ .

$$\begin{array}{ccc} (E, H) & & (\tilde{E}, \tilde{H}) \\ & \searrow \pi & \swarrow \tilde{\pi} \\ & M & \end{array}$$

In this setting another space which is important is the *correspondence space* which is the fiber product of the bundles  $E$  and  $\tilde{E}$ . The correspondence space projects over each of the T-dual spaces and has a natural 3-form on it:  $H - \tilde{H} = dA$ , where  $A = -\theta \wedge \tilde{\theta}$ .

$$\begin{array}{ccccc} & & (E \times_M \tilde{E}, p^*H - \tilde{p}^*\tilde{H}) & & (4.1) \\ & \swarrow p & & \searrow \tilde{p} & \\ (E, H) & & & & (\tilde{E}, \tilde{H}) \\ & \searrow \pi & & \swarrow \tilde{\pi} & \\ & & M & & \end{array}$$

We remark that although the space  $\tilde{E}$  is well defined from the data  $(E, H, \theta)$ , the same is not true about  $[\tilde{H}]$ , which is well defined up to an element in the ideal  $[F] \wedge H^1(\tilde{E}) \subset H^3(\tilde{E})$ .

**Example 4.1.** The Hopf fibration makes the 3-sphere,  $S^3$ , a principal  $S^1$  bundle over  $S^2$ . The curvature of this bundle is a volume form of  $S^2$ ,  $\sigma$ . So  $S^3$  equipped with the zero 3-form is T-dual to  $(S^2 \times S^1, \sigma \wedge \theta)$ . On the other hand, still considering the Hopf fibration, the 3-sphere endowed with the 3-form  $H = \theta \wedge \sigma$  is self T-dual.

**Example 4.2** (Lie Groups). Let  $(G, H)$  be a semi-simple Lie group with 3-form  $H(X, Y, Z) = K([X, Y], Z)$ , the Cartan form generating  $H^3(G, \mathbb{Z})$ , where  $K$  is the Killing form.

With a choice of an  $S^1 \subset G$ , we can think of  $G$  as a principal circle bundle. For  $X = \partial_\theta \in \mathfrak{g}$  tangent to  $S^1$  and of length  $-1$  according to the Killing form, a natural connection on  $G$  is given by  $-K(X, \cdot)$ . The curvature of this connection is given by

$$d(-K(X, \cdot))(Y, Z) = K(X, [Y, Z]) = H(X, Y, Z),$$

hence  $c_1$  and  $\tilde{c}_1$  are related by

$$c_1 = H(X, \cdot, \cdot) = X \lrcorner H = \tilde{c}_1.$$

Which shows that semi-simple Lie groups with the Cartan 3-form are self T-dual.

If  $E$  and  $\tilde{E}$  are T-dual spaces, we can define a map of invariant forms  $\tau : \Omega_{S^1}^\bullet(E) \longrightarrow \Omega_{S^1}^\bullet(\tilde{E})$  by  $\tau(\rho) = \tilde{p}_* e^A p^* \rho$ , where  $H - \tilde{H} = dA$ , or more explicitly

$$\tau(\rho) = \frac{1}{2\pi} \int_{S^1} e^A \rho. \quad (4.2)$$

If we decompose  $\rho = \rho_0 + \rho_1 \wedge \theta$ , with  $\rho_i$  pull back from  $M$ , then one can check that  $\tau$  is given by

$$\tau(\rho_0 + \rho_1 \wedge \theta) = \rho_1 + \theta \rho_0. \quad (4.3)$$

It is clear from (4.3) and that if we T-dualize twice and choose  $\theta = \tilde{\theta}$  for the second T-duality, we get  $(E, H)$  back and  $\tau^2 = \text{Id}$ .

The main theorem from [7] concerning us is:

**Theorem 4.3** (Bouwknegt–Evslin–Mathai [7]). *The map*

$$\tau : (\Omega_{S^1}^\bullet(E), d_H) \longrightarrow (\Omega_{S^1}^\bullet(\tilde{E}), d_{\tilde{H}})$$

*is an isomorphism of differential complexes.*

*Proof.* Given that  $\tau$  has an inverse, obtained by T-dualizing again, we only have to check that  $\tau$  preserves the differentials, i.e.,  $d_{\tilde{H}} \circ \tau = \tau \circ d_H$ . To obtain this relation we use equation (4.2):

$$\begin{aligned} d_{\tilde{H}}\tau(\rho) &= \frac{1}{2\pi} \int_{S^1} d_{\tilde{H}}(e^{-\theta\tilde{\theta}}\rho) \\ &= \frac{1}{2\pi} \int_{S^1} (H - \tilde{H})e^{-\theta\tilde{\theta}}\rho + e^{-\theta\tilde{\theta}}d\rho + \tilde{H}e^{-\theta\tilde{\theta}}\rho \\ &= \frac{1}{2\pi} \int_{S^1} He^{-\theta\tilde{\theta}}\rho + e^{-\theta\tilde{\theta}}d\rho \\ &= \tau(d_H\rho) \end{aligned}$$

□

*Remark:* If one considers  $\tau$  as a map of the complexes of differential forms (no invariance required), it will not be invertible. Nonetheless, every  $d_H$ -cohomology class has an invariant representative, hence  $\tau$  is a quasi-isomorphism.

### 4.1.1 Principal torus bundles

The construction of the T-dual described above can also be used to construct T-duals of principal torus bundles. What one has to do is just to split the torus into a product of circles and use the previous construction with “a circle at a time” (see [8]). However, this is only possible if

$$H(X, Y, \cdot) = 0 \quad \text{if } X, Y \text{ are vertical.} \quad (4.4)$$

Mathai and Rosenberg studied the case when (4.4) fails in [52]. There they propose that the T-dual is a field of noncommutative tori.

Another important difference between the circle bundle case and the torus bundle case is that in the former  $\tilde{E}$  is determined by  $E$  and  $[H]$  while in the latter this is not true [10]. One can see why this is the case if we recall that for circle bundles, even though  $\tilde{E}$  is well defined from  $E$  and  $[H]$ , the same is not true about  $[\tilde{H}]$ . So if one wants to T-dualize again, along a different circle direction, the topology of the next T-dual,  $\tilde{\tilde{E}}$ , will depend on  $\tilde{E}$  and  $[\tilde{H}]$  and hence is not well defined.

**Example 4.4.** A simple example to illustrate this fact is given by a 2-torus bundle with nonvanishing Chern classes but with  $[H] = 0$ . Taking the 3-form  $H = 0$  as a representative, a T-dual will be a flat torus bundle. Taking  $H = d(\theta_1 \wedge \theta_2) = c_1 \theta_2 - c_2 \theta_1$  as a representative of the zero cohomology class, a T-dual will be the torus bundle with (nonzero) Chern classes  $[c_1]$  and  $[-c_2]$ .

This fact leads us to define T-duality as a relation.

**Definition 4.5.** Let  $(E, H)$  and  $(\tilde{E}, \tilde{H})$  be principal  $n$ -torus bundles over a base  $M$ . We say that  $E$  and  $\tilde{E}$  are *T-dual* if on the correspondence space  $E \times_M \tilde{E}$  we have  $H - \tilde{H} = dA$ , where

$$[A]|_{T^n \times \tilde{T}^n} = \pm \sum \theta_i \wedge \tilde{\theta}_i \in H^2(T^n \times \tilde{T}^n)/H^2(T^n) \times H^2(\tilde{T}^n)$$

Clearly Theorem 4.3 still holds in this case with the same proof.

## 4.2 T-duality as a map of Courant algebroids

In this section we state our main result for T-dual circle bundles. The case of torus bundles can be dealt with similar techniques. We have seen that given two T-dual circle bundles we have a map of differential algebras  $\tau$  which is an isomorphism of the invariant differential exterior algebras. Now we introduce a map on invariant sections of generalized tangent spaces:

$$\varphi : T_{S^1} E \oplus T_{S^1}^* E \longrightarrow T_{S^1} \tilde{E} \oplus T_{S^1}^* \tilde{E}.$$

Any invariant section of  $TE \oplus TE^*$  can be written as  $X + f\partial_\theta + \xi + g\theta$ , where  $X$  is a horizontal vector and  $\xi$  is pull-back from the base. We define  $\varphi$  by:

$$\varphi(X + f\partial_\theta + \xi + g\theta) = X + g\partial_{\tilde{\theta}} + \xi + f\tilde{\theta}. \quad (4.5)$$

The relevance of this map comes from our main result.

**Theorem 4.6** (Cavalcanti–Gualtieri [18]). *The map  $\varphi$  defined in (4.5) is an orthogonal isomorphism of Courant algebroids and relates to  $\tau$  acting on invariant forms via*

$$\tau(V \cdot \rho) = \varphi(V) \cdot \tau(\rho). \quad (4.6)$$

*Proof.* It is obvious from equation (4.5) that  $\varphi$  is orthogonal with respect to the natural pairing. To prove equation (4.6) we observe that the map  $\tau$  defined in (4.2) can be interpreted as the right Clifford action of  $(\partial_\theta + \theta)$  followed by a relabeling  $\theta \mapsto \tilde{\theta}$ :

$$(\rho_0 + \rho_1 \wedge \theta) \cdot (\partial_\theta + \theta) = \rho_1 + \rho_0 \wedge \theta \mapsto \rho_1 + \rho_0 \wedge \tilde{\theta}.$$

In the same light, the map  $\varphi$  defined in (4.5) is just the reflexion on the orthogonal complement of  $\partial_\theta - \theta$  followed by relabeling  $\theta \mapsto \tilde{\theta}$ .

From this point of view, the result of this theorem is just about a commuting relation for right and left Clifford actions on forms. Given a form  $\rho$ , one can check that

$$\rho \cdot (\partial_\theta + \theta) = (-1)^{|\rho|+1} (\partial_\theta - \theta) \cdot \rho,$$

so for  $v \in TE \oplus T^*E$ , and  $\rho$  an even form we have

$$\begin{aligned} \tau(v \cdot \rho) &= (v \cdot \rho) \cdot (\partial_\theta + \theta) = (\partial_\theta - \theta) \cdot (v \cdot \rho) \\ &= (\partial_\theta - \theta) \cdot v \cdot (\partial_\theta - \theta) \cdot (-1)^{|\rho|} (\partial_\theta - \theta) \cdot \rho \\ &= \varphi(v) \cdot (\rho \cdot (\partial_\theta + \theta)) = \varphi(v) \cdot \tau(\rho). \end{aligned}$$

Finally, we have established that under the isomorphisms  $\varphi$  of Clifford algebras and  $\tau$  of Clifford modules,  $d_H$  corresponds to  $d_{\tilde{H}}$ , hence the induced brackets (according to equation (1.9)) are the same.  $\square$

*Remark:* As  $E$  is the total space of a circle bundle, its invariant tangent bundle sits in the Atiyah sequence:

$$0 \longrightarrow 1 = T_1 S^1 \longrightarrow T_{S^1} E \longrightarrow TM \longrightarrow 0$$

or, taking duals,

$$0 \longrightarrow T^*M \longrightarrow T_{S^1}^* E \longrightarrow T_1^* S^1 = 1^* \longrightarrow 0.$$

The choice of a connection on  $E$  induces a splitting of the sequences above and an isomorphism

$$T_{S^1} E \oplus T_{S^1}^* E \cong TM \oplus T^*M \oplus 1 \oplus 1^*,$$



The argument also applies to  $\tilde{E}$ :

$$T_{S^1}\tilde{E} \oplus T_{S^1}^*\tilde{E} \cong TM \oplus T^*M \oplus 1 \oplus 1^*.$$

The map  $\varphi$  can be seen in this light as the permutation of the terms 1 and  $1^*$ . This is Ben-Bassat's starting point for the study of mirror symmetry and generalized complex structures in [4, 5].

Since the Courant algebroids  $(T_{S^1}E \oplus T_{S^1}^*E, \llbracket \cdot, \cdot \rrbracket_H)$  and  $(T_{S^1}\tilde{E} \oplus T_{S^1}^*\tilde{E}, \llbracket \cdot, \cdot \rrbracket_{\tilde{H}})$  are isomorphic, according to Theorem 4.6, we see that any invariant structure on  $(TE \oplus T^*E, \llbracket \cdot, \cdot \rrbracket_H)$  defined in terms of the Courant bracket and natural pairing correspond to a similar structure on  $(T\tilde{E} \oplus T^*\tilde{E}, \llbracket \cdot, \cdot \rrbracket_{\tilde{H}})$ .

**Theorem 4.7** (Cavalcanti–Gualtieri [18]). *Any invariant Dirac, generalized complex, generalized Kähler on  $(TE \oplus T^*E, \llbracket \cdot, \cdot \rrbracket_H)$  is transformed into a similar one via  $\varphi$ .*

**Exercise 4.8.** What happens with the generalized Kähler structure on Lie groups described in Example 2.10 under T-duality?

The decomposition of  $\wedge^*T_{\mathbb{C}}^*M$  into subbundles  $U^k$  is also preserved from T-duality.

**Corollary 4.9.** *If two generalized complex manifolds  $(E, \mathcal{J}_1)$  and  $(\tilde{E}, \mathcal{J}_2)$  correspond via T-duality, then  $\tau(\mathcal{U}_E^k) = \mathcal{U}_{\tilde{E}}^k$  and also*

$$\tau(\partial_E\psi) = -\partial_{\tilde{E}}\tau(\psi) \quad \tau(\bar{\partial}_E\psi) = -\bar{\partial}_{\tilde{E}}\tau(\psi).$$

*Proof.* The T-dual generalized complex structure in  $\tilde{E}$  is determined by  $\tilde{L} = \varphi(L)$ , where  $L$  is the  $+i$ -eigenspace of the generalized complex structure on  $E$ . Since  $\varphi$  is real,  $\tilde{\bar{L}} = \varphi(\bar{L})$ , and hence

$$\mathcal{U}_{\tilde{E}}^{n-k} = \Omega^k(\tilde{\bar{L}}) \cdot \tau(\rho) = \tau(\Omega^k(\bar{L}) \cdot \rho) = \tau(\mathcal{U}_E^k).$$

Finally, if  $\alpha \in \mathcal{U}^k$ , then

$$\partial_{\tilde{E}}\tau(\alpha) - \bar{\partial}_{\tilde{E}}\tau(\alpha) = d_{\tilde{H}}\tau(\alpha) = -\tau(d_H\alpha) = -\tau(\partial_E\alpha) + \tau(\bar{\partial}_E\alpha).$$

Since  $\tau(\mathcal{U}^k) = \mathcal{U}_{\tilde{E}}^k$ , we obtain the identities for the operators  $\partial_{\tilde{E}}$  and  $\bar{\partial}_{\tilde{E}}$ .  $\square$

**Example 4.10** (Change of type of generalized complex structures). As even and odd forms get swapped with T-duality along a circle, the type of a generalized complex structure is not preserved. However, it can only change, at a point, by  $\pm 1$ . Indeed, if  $\rho = e^{B+i\omega}\Omega$  is an invariant form determining a generalized complex structure there are two possibilities: If  $\Omega$  is a pull back from the base, the type will increase by 1, otherwise will decrease by 1.

For a principal  $n$ -torus bundle, the rule is not so simple. If we let  $T^n$  be the fiber,  $\rho = e^{B+i\omega}\Omega$  be a local trivialization of the canonical bundle and define

$$l = \max\{i : \wedge^i TT \cdot \Omega \neq 0\}$$

and

$$r = \text{rank}(\omega|_V), \text{ where } V = \text{Ann}(\Omega) \cap TT,$$

then the type,  $\tilde{t}$  of the T-dual structure relates to the type,  $t$ , of the original structure by

$$\tilde{t} = t + n - 2l - r. \quad (4.7)$$

The following table summarizes different ways the type changes for generalized complex structures in  $E^{2n}$  induced by complex and symplectic structures if the fibers are  $n$ -tori of some special types:

Structure on $E$	Fibers of $E$	Structure on $\tilde{E}$	Fibers of $\tilde{E}$
Complex	Complex	Complex	Complex
Complex	Real	Symplectic	Lagrangian
Symplectic	Symplectic	Symplectic	Symplectic
Symplectic	Lagrangian	Complex	Real

Table 1: Change of type of generalized complex structures under T-duality according to the type of fiber.

**Example 4.11** (Hopf surfaces). Given two complex numbers  $a_1$  and  $a_2$ , with  $|a_1|, |a_2| > 1$ , the quotient of  $\mathbb{C}^2$  by the action  $(z_1, z_2) \mapsto (a_1 z_1, a_2 z_2)$  is a primary Hopf surface (with the induced complex structure). Of all primary Hopf surfaces, these are the only ones admitting a  $T^2$  action preserving the complex structure (see [3]). If  $a_1 = a_2$ , the orbits of the 2-torus action are elliptic surfaces and hence, according to Example 4.10, the T-dual will still be a complex

manifold. If  $a_1 \neq a_2$ , then the orbits of the torus action are real except for the orbits passing through  $(1, 0)$  and  $(0, 1)$ , which are elliptic. In this case, the T-dual will be generically symplectic except for the two special fibers corresponding to the elliptic curves, where there is type change. This example also shows that even if the initial structure on  $E$  has constant type, the same does not need to be true in the T-dual.

**Example 4.12** (Mirror symmetry of Betti numbers). Consider the case of the mirror of a Calabi-Yau manifold along a special Lagrangian fibration. We have seen that the bundles  $U_{\omega, J}^k$  induced by both the complex and symplectic structure are preserved by T-duality. Hence  $U^{p, q} = U_{\omega}^p \cap U_J^q$  is also preserved, but,  $U^{p, q}$  will be associated in the mirror to  $U_J^p \cap U_{\omega}^q$ , as complex and symplectic structure get swapped. Finally, as remarked Chapter 2, example 2.2, we have an isomorphism between  $\Omega^{p, q}$  and  $\mathcal{U}^{n-p-q, p-q}$ . Making these identifications, we have

$$\Omega^{p, q}(E) \cong \mathcal{U}^{n-p-q, p-q}(E) \cong \tilde{\mathcal{U}}^{n-p-q, p-q}(\tilde{E}) \cong \Omega^{n-p, q}(\tilde{E}).$$

Which, in cohomology, gives the usual ‘mirror symmetry’ of the Hodge diamond.

### 4.2.1 The metric and the Buscher rules

Another geometric structure that can be transported via T-duality is the generalized metric. Assume that a principal circle bundle  $E$  is endowed with an invariant generalized metric  $\mathcal{G}$ . Then, since  $\varphi$  is orthogonal,  $\tilde{\mathcal{G}} = \varphi \mathcal{G} \varphi^{-1}$  is a generalized metric on  $\tilde{E}$  and with these metrics  $\varphi$  is an isometry between  $TE \oplus T^*E$  and  $T\tilde{E} \oplus T^*\tilde{E}$ .

Since a generalized metric in a split Courant algebroid is defined by a metric and a 2-form,  $\mathcal{G}$  is equivalent to an invariant metric  $g$  and an invariant 2-form  $b$  which we can write as

$$\begin{aligned} g &= g_0 \theta \odot \theta + g_1 \odot \theta + g_2 \\ b &= b_1 \wedge \theta + b_2. \end{aligned}$$

If one wants to determine the corresponding metric  $\tilde{g}$  and 2-form  $b$  on  $\tilde{E}$  we just have to recall that the 1-eigenspace of  $\tilde{\mathcal{G}}$ ,  $\tilde{C}_+ = \varphi(C_+)$ ,

is the graph of  $\tilde{g} + \tilde{b}$ . One can check that  $\tilde{C}_+$  is the graph of:

$$\begin{aligned}\tilde{g} &= \frac{1}{g_0} \tilde{\theta} \odot \tilde{\theta} - \frac{b_1}{g_0} \odot \tilde{\theta} + g_2 + \frac{b_1 \odot b_1 - g_1 \odot g_1}{g_0} \\ \tilde{b} &= -\frac{g_1}{g_0} \wedge \tilde{\theta} + b_2 + \frac{g_1 \wedge b_1}{g_0}\end{aligned}\tag{4.8}$$

Of course, in the generalized Kähler case, this is how the  $g$  and  $b$  induced by the structure transform. These equations, however, are not new. They had been encountered before by the physicists [13, 14], independently of generalized complex geometry and are called *Buscher rules!*

### 4.2.2 The bihermitian structure

The choice of a generalized metric  $(g, b)$  gives us two orthogonal spaces

$$C_{\pm} = \{X + b(X, \cdot) \pm g(X, \cdot) : X \in TM\},$$

and the projections  $\pi_{\pm} : C_{\pm} \rightarrow TM$  are isomorphisms. Hence, any endomorphism  $A \in \text{End}(TM)$  induces endomorphisms  $A_{\pm}$  on  $C_{\pm}$ . Using the map  $\varphi$  we can transport this structure to a T-dual:

$$\begin{array}{ccc} A_+ \in \text{End}(C_+) & \xrightarrow{\varphi} & \tilde{A}_+ \in \text{End}(\tilde{C}_+) \\ \pi_+ \downarrow & & \tilde{\pi}_+ \downarrow \\ A \in \text{End}(TE) & & \tilde{A}_{\pm} \in \text{End}(T\tilde{E}) \\ \pi_- \uparrow & & \tilde{\pi}_- \uparrow \\ A_- \in \text{End}(C_-) & \xrightarrow{\varphi} & \tilde{A}_- \in \text{End}(\tilde{C}_-)\end{array}$$

As we are using the generalized metric to transport  $A$  and the maps  $\pi_{\pm}$  and  $\varphi$  are orthogonal, the properties shared by  $A$  and  $A_{\pm}$  will be metric related ones, e.g., self-adjointness, skew-adjointness and orthogonality. In the generalized Kähler case, it is clear that if we transport  $J_{\pm}$  via  $C_{\pm}$  we obtain the corresponding complex structures

of the induced generalized Kähler structure in the dual:

$$\tilde{J}_\pm = \tilde{\pi}_\pm \varphi \pi_\pm^{-1} J_\pm (\tilde{\pi}_\pm \varphi \pi_\pm^{-1})^{-1}.$$

In the case of a metric connexion,  $\theta = g(\partial_\theta, \cdot)/g(\partial_\theta, \partial_\theta)$ , we can give a very concrete description of  $\tilde{J}_\pm$ . We start describing the maps  $\tilde{\pi}_\pm \varphi \pi_\pm^{-1}$ . If  $V$  is orthogonal to  $\partial_\theta$ , then  $g_1(V) = 0$  and

$$\begin{aligned} \tilde{\pi}_\pm \varphi \pi_\pm^{-1}(V) &= \tilde{\pi}_\pm \varphi(V + b_1(V)\theta + b_2(V) \pm g_2(V, \cdot)) \\ &= \tilde{\pi}_\pm(V + b_1(V) \frac{\partial}{\partial \tilde{\theta}} + b_2(V) \pm g_2(V, \cdot)) \\ &= V + b_1(V) \frac{\partial}{\partial \tilde{\theta}}. \end{aligned}$$

And for  $\partial_\theta$  we have

$$\tilde{\pi}_\pm \varphi \pi_\pm^{-1}(\partial_\theta) = \tilde{\pi}_\pm \varphi(\partial_\theta + b_1 \pm (\frac{1}{g_0}\theta + g_1)) = \tilde{\pi}_\pm(\frac{1}{g_0}\partial_{\tilde{\theta}} + \tilde{\theta}) = \pm \frac{1}{g_0}\partial_{\tilde{\theta}}.$$

*Remark:* The T-dual connection is *not* the metric connection for the T-dual metric. This is particularly clear in this case, as the vector  $\tilde{\pi}_\pm \varphi \pi_\pm^{-1}(V) = V + b_1(V)\partial_{\tilde{\theta}}$ , although not horizontal for the T-dual connection, is perpendicular to  $\partial_{\tilde{\theta}}$  according to the dual metric. This means that if we use the metric connections of both sides, the map  $\tilde{\pi}_\pm \varphi \pi_\pm^{-1}$  is the identity from the orthogonal complement of  $\partial_\theta$  to the orthogonal complement of  $\partial_{\tilde{\theta}}$ .

Now, if we let  $V_\pm$  be the orthogonal complement to  $\text{span}\{\partial_\theta, J_\pm \partial_\theta\}$  we can describe  $\tilde{J}_\pm$  by

$$\tilde{J}_\pm w = \begin{cases} J_\pm w, & \text{if } w \in V_\pm \\ \pm \frac{1}{g_0} J_\pm \partial_\theta & \text{if } w = \partial_\theta \\ \mp g_0 \partial_{\tilde{\theta}} & \text{if } w = J_\pm \partial_\theta \end{cases} \quad (4.9)$$

Therefore, if we identify  $\partial_\theta$  with  $\partial_{\tilde{\theta}}$  and their orthogonal complements with each other via  $TM$ ,  $\tilde{J}_+$  is essentially the same as  $J_+$ , but stretched in the directions of  $\partial_\theta$  and  $J_+ \partial_\theta$  by  $g_0$ , while  $\tilde{J}_-$  is  $J_-$  conjugated and stretched in those directions. In particular,  $J_+$  and  $\tilde{J}_+$  determine the same orientation while  $\tilde{J}_-$  and  $J_-$  determine reverse orientations.

### 4.3 Reduction and T-duality

Now, let  $(E, H)$  and  $(\tilde{E}, \tilde{H})$  be T-dual spaces and consider the correspondence space  $E \times_M \tilde{E}$  with the 3-form  $p^*H$ :

$$\begin{array}{ccc} & (E \times_M \tilde{E}, p^*H) & \\ p \swarrow & & \searrow \tilde{p} \\ E & & \tilde{E} \end{array}$$

There are two circle actions on this space with associated Lie algebra maps  $\psi_i : \mathbb{R} \rightarrow \Gamma(TM)$ ,  $\psi_1(1) = \partial_\theta$  and  $\psi_2(1) = \partial_{\tilde{\theta}}$ . Since  $H$  is basic with respect to the action of  $\partial_{\tilde{\theta}}$ , we can lift the action induced by  $\psi_2$  and form the corresponding reduced algebroid over  $E = M/S^1$ , which is just  $(TE + T^*E, [, ]_H, \langle \cdot, \cdot \rangle)$ .

On the other hand,  $H$  has an equivariantly closed extension with respect to the action of  $\partial_\theta$ , since  $i\partial_\theta H = d\tilde{\theta}$ , so we can lift the action of  $\partial_\theta$  as  $\rho_1(1) = \partial_\theta - \tilde{\theta}$ . As in Example 3.14, the connection  $\theta$  allows us to choose a natural splitting for the reduced algebroid over  $\tilde{E}$ , which according to (3.7) has curvature  $\tilde{H} = H - d(\theta \wedge \tilde{\theta}) = (d\theta) \wedge \tilde{\theta} + h$ , hence we have the following

$$\begin{array}{ccc} & (E \times_M \tilde{E}, p^*H) & \\ / \partial_{\tilde{\theta}} \swarrow & & \searrow / \frac{\partial}{\partial \tilde{\theta}} - \tilde{\theta} \\ (E, H) & & (\tilde{E}, \tilde{H}) \end{array}$$

Observe that for the first reduction we had  $K_1 = \{\partial_{\tilde{\theta}}\}$  and for the second reduction we had  $K_2 = \{\partial_\theta + \tilde{\theta}\}$  and the natural pairing gives a nondegenerate pairing between these two spaces.

**Theorem 4.13** (Cavalcanti–Gualtieri [18], Hu [39]). *If two principal torus bundles over a common base  $(E, H)$  and  $(\tilde{E}, \tilde{H})$  are T-dual to each other then they can be obtained as reduced spaces from a common space  $(M, \mathcal{H})$  by two torus actions.*

*If  $K_1$  and  $K_2$  are the vector bundles generated by the lifts of each of these actions to the Courant algebroid  $(TM + T^*M, [, ]_{\mathcal{H}}, \langle \cdot, \cdot \rangle)$ , then  $K_1$  and  $K_2$  are isotropic and the natural pairing is nondegenerate in  $K_1 \times K_2 \rightarrow \mathbb{R}$ .*

Finally, we observe that reducing  $(TM + T^*M, [, ]_{\mathcal{H}}, \langle \cdot, \cdot \rangle)$  by the full  $T^{2n}$  action renders a Courant algebroid over the common base  $M$ . The rank of this Courant algebroid is the same as the rank of the reduced algebroids over either  $E$  or  $\tilde{E}$  and it can be geometrically interpreted in two different ways: invariant sections of  $TE + T^*E$  or invariant sections  $T\tilde{E} + T^*\tilde{E}$ . Of course the algebroid itself does not depend on the particular interpretation, hence  $(TE + T^*E)_{T^n}$  and  $(T\tilde{E} + T^*\tilde{E})_{T^n}$  are isomorphic as Courant algebroids over  $M$ , which is precisely the result of Theorem 4.6





# Bibliography

- [1] V. APOSTOLOV, P. GAUDUCHON, AND G. GRANTCHAROV. Bi-Hermitian structures on complex surfaces. *Proc. London Math. Soc.* (3), 79(2):414–428, 1999.
- [2] V. APOSTOLOV AND M. GUALTIERI. Generalized Kähler manifolds with split tangent bundle, 2006. math.DG/0605342, to appear in *Comm. Math. Phys.*
- [3] W. BARTH, C. PETERS, AND A. VAN DE VEN. *Compact complex surfaces*, volume 4 of *Ergebnisse der Mathematik und ihrer Grenzgebiete (3) [Results in Mathematics and Related Areas (3)]*. Springer-Verlag, Berlin, 1984.
- [4] O. BEN-BASSAT. Mirror symmetry and generalized complex manifolds. I. The transform on vector bundles, spinors, and branes. *J. Geom. Phys.*, 56(4):533–558, 2006.
- [5] O. BEN-BASSAT. Mirror symmetry and generalized complex manifolds. II. Integrability and the transform for torus bundles. *J. Geom. Phys.*, 56(7):1096–1115, 2006.
- [6] O. BEN-BASSAT AND M. BOYARCHENKO. Submanifolds of generalized complex manifolds. *J. Symplectic Geom.*, 2(3):309–355, 2004.
- [7] P. BOUWKNEGT, J. EVSLIN, AND V. MATHAI.  $T$ -duality: topology change from  $H$ -flux. *Comm. Math. Phys.*, 249(2):383–415, 2004. hep-th/0306062.

- [8] P. BOUWKNEGT, K. HANNABUSS, AND V. MATHAI.  $T$ -duality for principal torus bundles. *J. High Energy Phys.*, 3:018, 10 pp. (electronic), 2004. hep-th/0312284.
- [9] J.-L. BRYLINSKI. A differential complex for Poisson manifolds. *J. Differential Geom.*, 28(1):93–114, 1988.
- [10] U. BUNKE AND T. SCHICK. On the topology of  $T$ -duality. *Rev. Math. Phys.*, 17(1):77–112, 2005.
- [11] H. BURSZTYN, G. R. CAVALCANTI, AND M. GUALTIERI. Reduction of Courant algebroids and generalized complex structures. math.DG/0509640. To appear in *Adv. Math.*, 2005.
- [12] H. BURSZTYN AND O. RADKO. Gauge equivalence of Dirac structures and symplectic groupoids. *Ann. Inst. Fourier (Grenoble)*, 53(1):309–337, 2003.
- [13] T. H. BUSCHER. A symmetry of the string background field equations. *Phys. Lett. B*, 194(1):59–62, 1987.
- [14] T. H. BUSCHER. Path-integral derivation of quantum duality in nonlinear sigma-models. *Phys. Lett. B*, 201(4):466–472, 1988.
- [15] G. R. CAVALCANTI. Formality in generalized Kähler geometry. math.DG/060396, to appear in *Topology Appl.*, 2006.
- [16] G. R. CAVALCANTI. The decomposition of forms and cohomology of generalized complex manifolds. *J. Geom. Phys.*, 57:121–132, 2007.
- [17] G. R. CAVALCANTI. The Lefschetz property, formality and blowing up in symplectic geometry. *Trans. Amer. Math. Soc.*, 359:333–348, 2007.
- [18] G. R. CAVALCANTI AND M. GUALTIERI.  $T$ -duality and generalized complex geometry. Still in preparation...
- [19] G. R. CAVALCANTI AND M. GUALTIERI. Generalized complex structures on nilmanifolds. *J. Symplectic Geom.*, 2(3):393–410, 2004.

- [20] G. R. CAVALCANTI AND M. GUALTIERI. A surgery for generalized complex structures on 4-manifolds, 2006. math.DG/0602333. To appear in *J. Differential Geom.*
- [21] C. CHEVALLEY. *The algebraic theory of spinors and Clifford algebras*. Springer-Verlag, Berlin, 1997. Collected works. Vol. 2, Edited and with a foreword by Pierre Cartier and Catherine Chevalley, With a postface by J.-P. Bourguignon.
- [22] T. COURANT AND A. WEINSTEIN. Beyond Poisson structures. In *Action hamiltoniennes de groupes. Troisième théorème de Lie (Lyon, 1986)*, volume 27 of *Travaux en Cours*, pages 39–49. Hermann, Paris, 1988.
- [23] T. J. COURANT. Dirac manifolds. *Trans. Amer. Math. Soc.*, 319(2):631–661, 1990.
- [24] L. C. DE ANDRES, M. L. BARBERIS, I. DOTTI, AND M. FERNANDEZ. Hermitian structures on cotangent bundles of four dimensional solvable lie groups, 2006. math.DG/0604608.
- [25] P. DELIGNE, P. GRIFFITHS, J. MORGAN, AND D. SULLIVAN. Real homotopy theory of Kähler manifolds. *Invent. Math.*, 29(3):245–274, 1975.
- [26] M. FERNÁNDEZ, R. IBÁÑEZ, AND M. DE LEÓN. On a Brylinski conjecture for compact symplectic manifolds. In *Quaternionic structures in mathematics and physics (Trieste, 1994)*, pages 119–126 (electronic). Int. Sch. Adv. Stud. (SISSA), Trieste, 1998.
- [27] S. FIDANZA, R. MINASIAN, AND A. TOMASIELLO. Mirror symmetric SU(3)-structure manifolds with NS fluxes. *Comm. Math. Phys.*, 254(2):401–423, 2005. hep-th/0311122.
- [28] J. FIGUEROA-O’FARRILL AND N. MOHAMMEDI. Gauging the Wess-Zumino term of a sigma model with boundary. *J. High Energy Phys.*, (8):086, 22 pp. (electronic), 2005.
- [29] J. M. FIGUEROA-O’FARRILL AND S. STANCIU. Gauged Wess-Zumino terms and equivariant cohomology. *Phys. Lett. B*, 341(2):153–159, 1994.

- [30] S. J. GATES, JR., C. M. HULL, AND M. ROČEK. Twisted multiplets and new supersymmetric nonlinear  $\sigma$ -models. *Nuclear Phys. B*, 248(1):157–186, 1984.
- [31] R. E. GOMPF AND T. S. MROWKA. Irreducible 4-manifolds need not be complex. *Ann. of Math. (2)*, 138(1):61–111, 1993.
- [32] M. GOZE AND Y. KHAKIMDJANOV. *Nilpotent Lie algebras*, volume 361 of *Mathematics and its Applications*. Kluwer Academic Publishers Group, Dordrecht, 1996.
- [33] M. GRAÑA, R. MINASIAN, M. PETRINI, AND A. TOMASIELLO. Supersymmetric backgrounds from generalized Calabi-Yau manifolds. *J. High Energy Phys.*, (8):046, 34 pp. (electronic), 2004.
- [34] M. GUALTIERI. *Generalized Complex Geometry*. D.Phil. thesis, Oxford University, 2003. math.DG/0401221.
- [35] M. GUALTIERI. Generalized geometry and the Hodge decomposition. math.DG/0409093, 2004.
- [36] S. GURRIERI, J. LOUIS, A. MICU, AND D. WALDRAM. Mirror symmetry in generalized Calabi-Yau compactifications. *Nuclear Phys. B*, 654(1-2):61–113, 2003. hep-th/0211102.
- [37] N. HITCHIN. Generalized Calabi-Yau manifolds. *Q. J. Math.*, 54(3):281–308, 2003.
- [38] S. HU. Hamiltonian symmetries and reduction in generalized geometry, 2005.
- [39] S. HU. Reduction and duality in generalized geometry, 2005. math.DG/0512634.
- [40] C. M. HULL AND B. SPENCE. The gauged nonlinear sigma model with Wess-Zumino term. *Phys. Lett. B*, 232(2):204–210, 1989.
- [41] R. IBÁÑEZ, Y. RUDYAK, A. TRALLE, AND L. UGARTE. Symplectically harmonic cohomology of nilmanifolds. In *Symplectic and contact topology: interactions and perspectives (Toronto, ON/Montreal, QC, 2001)*, volume 35 of *Fields Inst. Commun.*, pages 117–130. Amer. Math. Soc., Providence, RI, 2003.

- [42] A. KAPUSTIN AND D. ORLOV. Remarks on A-branes, mirror symmetry, and the Fukaya category. *J. Geom. Phys.*, 48(1):84–99, 2003.
- [43] M. K. KINYON AND A. WEINSTEIN. Leibniz algebras, Courant algebroids, and multiplications on reductive homogeneous spaces. *Amer. J. Math.*, 123(3):525–550, 2001.
- [44] S. KOBAYASHI AND K. NOMIZU. *Foundations of differential geometry. Vol. II.* Wiley Classics Library. John Wiley & Sons Inc., New York, 1996. Reprint of the 1969 original, A Wiley-Interscience Publication.
- [45] K. KODAIRA. On the structure of compact complex analytic surfaces. I. *Amer. J. Math.*, 86:751–798, 1964.
- [46] J.-L. KOSZUL. Crochet de Schouten-Nijenhuis et cohomologie. *Astérisque*, Numero Hors Serie:257–271, 1985. The mathematical heritage of Élie Cartan (Lyon, 1984).
- [47] Y. LI. On deformations of generalized complex structures: the generalized Calabi-Yau case, 2005. hep-th/0508030.
- [48] Y. LIN AND S. TOLMAN. Symmetries in generalized Kähler geometry. math.DG/0509069, to appear in *Comm. Math. Phys.*, 2005.
- [49] U. LINDSTRÖM, R. MINASIAN, A. TOMASIELLO, AND M. ZABZINE. Generalized complex manifolds and supersymmetry. *Comm. Math. Phys.*, 257(1):235–256, 2005.
- [50] U. LINDSTRÖM, M. ROČEK, R. VON UNGE, AND M. ZABZINE. Generalized Kähler geometry and manifest  $N = (2, 2)$  supersymmetric nonlinear sigma-models. *J. High Energy Phys.*, (7):067, 18 pp. (electronic), 2005.
- [51] Z.-J. LIU, A. WEINSTEIN, AND P. XU. Manin triples for Lie bialgebroids. *J. Differential Geom.*, 45(3):547–574, 1997.
- [52] V. MATHAI AND J. ROSENBERG.  $T$ -duality for torus bundles with  $H$ -fluxes via noncommutative topology. *Comm. Math. Phys.*, 253(3):705–721, 2005. hep-th/0401168.

- [53] O. MATHIEU. Harmonic cohomology classes of symplectic manifolds. *Comment. Math. Helv.*, 70(1):1–9, 1995.
- [54] D. MCDUFF AND D. SALAMON. *Introduction to symplectic topology*. Oxford Mathematical Monographs. The Clarendon Press Oxford University Press, New York, second edition, 1998.
- [55] S. A. MERKULOV. Formality of canonical symplectic complexes and Frobenius manifolds. *Internat. Math. Res. Notices*, 14:727–733, 1998.
- [56] S. MUKAI. Symplectic structure of the moduli space of sheaves on an abelian or  $K3$  surface. *Invent. Math.*, 77(1):101–116, 1984.
- [57] S. M. SALAMON. Complex structures on nilpotent Lie algebras. *J. Pure Appl. Algebra*, 157(2-3):311–333, 2001.
- [58] H. SAMELSON. A class of complex-analytic manifolds. *Portugaliae Math.*, 12:129–132, 1953.
- [59] P. ŠEVERA AND A. WEINSTEIN. Poisson geometry with a 3-form background. *Progr. Theoret. Phys. Suppl.*, 144:145–154, 2001. Noncommutative geometry and string theory (Yokohama, 2001).
- [60] M. STIENON AND P. XU. Reduction of generalized complex structures. math.DG/0509393, 2005.
- [61] D. SULLIVAN. Infinitesimal computations in topology. *Inst. Hautes Études Sci. Publ. Math.*, 47:269–331 (1978), 1977.
- [62] C. H. TAUBES. The Seiberg-Witten invariants and symplectic forms. *Math. Res. Lett.*, 1(6):809–822, 1994.
- [63] W. P. THURSTON. Some simple examples of symplectic manifolds. *Proc. Amer. Math. Soc.*, 55(2):467–468, 1976.
- [64] H.-C. WANG. Complex parallisable manifolds. *Proc. Amer. Math. Soc.*, 5:771–776, 1954.
- [65] A. WEINSTEIN. Symplectic manifolds and their Lagrangian submanifolds. *Advances in Math.*, 6:329–346 (1971), 1971.

- [66] A. WEINSTEIN. Omni-Lie algebras. *Sūrikaisekikenyūsho Kōkyūroku*, 1176:95–102, 2000. Microlocal analysis of the Schrödinger equation and related topics (Japanese) (Kyoto, 1999).
- [67] F. WITT. *Special metric structures and closed forms*. D.Phil. thesis, Oxford University, 2004. math.DG/0502443.
- [68] D. YAN. Hodge structure on symplectic manifolds. *Adv. Math.*, 120(1):143–154, 1996.
- [69] R. ZUCCHINI. A sigma model field theoretic realization of Hitchin’s generalized complex geometry. *J. High Energy Phys.*, (11):045, 21 pp. (electronic) (2005), 2004.
- [70] R. ZUCCHINI. Generalized complex geometry, generalized branes and the Hitchin sigma model. *J. High Energy Phys.*, 3:022, 33 pp. (electronic), 2005.