

ROBUSTLY EXPANSIVE CODIMENSION-ONE HOMOCLINIC CLASSES ARE HYPERBOLIC

M. J. PACIFICO, E. R. PUJALS, M. SAMBARINO, J. L. VIEITEZ

ABSTRACT. We shall prove that C^1 -robustly expansive codimension-one homoclinic classes are hyperbolic.

1. INTRODUCTION

Let M be a d -dimensional manifold and $\text{Diff}^1(M)$ be the set of C^1 diffeomorphisms f on M endowed with the C^1 topology. Let p be a hyperbolic periodic point of f and $H(p)$ be its homoclinic class. The diffeomorphism f is α -expansive in $H(p)$ if $\text{dist}(f^n(x), f^n(y)) \leq \alpha$ for all $n \in \mathbb{Z}$, with $x, y \in H(p)$, implies $x = y$. It is well known that hyperbolicity implies α -expansiveness for some $\alpha > 0$. But expansiveness alone does not guarantee hyperbolicity, even when one is dealing with a codimension one expansive homoclinic class, as can be seen in [PPV, Section 2]. We note that there are even examples of expansive codimension one homoclinic classes such all of its periodic orbits are hyperbolic that are not hyperbolic.

To see this consider a Smale horse-shoe H in a plane and Λ a non trivial minimal subset of H . In a complementary direction multiply H by a non uniform contraction $\lambda(w)$, depending on the distance from w to Λ , and in such way that $\lambda(w) = 1$ for $w \in \Lambda$. The resulting homoclinic class is expansive, has all periodic points hyperbolic but it is not hyperbolic.

Then we assume that expansiveness holds in a C^1 -neighborhood of the homoclinic class, that is, for all diffeomorphism g C^1 near f the homoclinic class $H(p_g)$ of the continuation p_g of p is α -expansive. In [PPV] it was proved that robustly expansive homoclinic classes of a three dimensional manifold are generically hyperbolic. We generalize this result in two ways. First we drop the assumption $\dim(M) = 3$ and get that robustly expansive codimension-one homoclinic classes have a codimension-one dominated splitting. Second we prove that robustly expansive codimension-one homoclinic classes with a dominated splitting are hyperbolic.

The main step to obtain this result is to prove that center unstable manifolds for all point x in the homoclinic class $H(p)$ are true unstable manifolds, that is, center unstable manifolds are dynamically defined. To achieve this it is enough to get this property for periodic points homoclinically related to p . For this we use a result of Sambarino that controls the behavior of periodic points homoclinically related to p . Once this is settled, under the hypothesis that the dominated splitting $E \oplus F$ is not hyperbolic, the fact that center unstable manifolds are dynamically defined allows us to obtain a hyperbolic periodic point q with arbitrarily large period as near $H(p)$ as we wish, and with arbitrarily small rate of contraction along the E -direction. Then we prove that

Date: January 16, 2006.

M.J.Pacifico and E.R. Pujals were partially supported by CNPq-Brazil/Faperj-Brazil/Pronex Dyn. Systems.

such hyperbolic periodic points are in fact in $H(p)$, contradicting uniform E -contraction in the period for periodic points in $H(p)$. As in [Ma2] this implies that the sub-bundle F is uniformly expanding, and then $E \oplus F$ is hyperbolic.

REFERENCES

- [Do] A. DOLD, *Lectures on algebraic topology*, Classics in Mathematics. Springer-Verlag, Berlin,, **1995** ISBN: 3-540-58660-1.
- [Fr] J. FRANKS, *Necessary conditions for stability if diffeomorphisms*, Trans. Amer. Math. Soc., **158** (1971), p. 301-308.
- [Ha] S. HAYASHI, *Connecting invariant manifolds and the solution of the C^1 -stability and Ω -stability conjecture for flows*, Ann. of Math. (2), **Vol. 145 N° 1** (1997), p. 81-137.
- [Li] S.T.LIAO, *On the stability conjecture*, Chinese Ann. Math., **1** (1980), p. 9-30.
- [Ma2] R. MAÑÉ, *A Proof of the C^1 Stability Conjecture*, Inst. Hautes Études Sci. Publ. Math., **66** (1988), p. 161-210.
- [Ma3] R. MAÑÉ, *An ergodic closing lemma*, Annals of Mathematics, **116** (1982), p. 503-540.
- [PM] J. PALIS AND W. DE MELO, *Geometric Theory of Dynamical Systems*, ., **Springer Verlag** 1982..
- [PI1] V. A. PLISS, *Analysis of the necessity of the conditions of Smale and Robbin for structural stability of periodic systems of differential equations*, Diff. Uravnenija, **8** (1972), p. 972-983.
- [PI2] V. A. PLISS, *On a conjecture due to Smale*, Diff. Uravnenija, **8** (1972), p. 268-282.
- [PS] E. PUJALS, M. SAMBARINO, *Homoclinic tangencies and hyperbolicity for surface diffeomorphisms*, Annals of Mathematics, **151** (2000), p. 961-1023.
- [PS2] E. PUJALS, M. SAMBARINO, *On the Dynamics of Dominated Splitting*, Preprint, (2002), p. -.
- [PPV] M. J. PACIFICO, E. R. PUJALS, J. L. VIEITEZ, *Robust expansive homoclinic classes*, Ergodic Theory and Dynamical Systems, **25** (2005), p. 271-300.
- [SV] SAMBARINO, M.; VIEITEZ, J., *On C^1 -persistently Expansive Homoclinic Classes*, Accepted for publication in Discrete and Continuous Dynamical Systems, (2005), p. -
- [1] [We2]WL2 L. WEN, *Heteroclinic Cycles and Homoclinic Closures for Generic Diffeomorphisms*, Preprint, (2002), p. -.