

# ROBUSTLY EXPANSIVE CODIMENSION-ONE HOMOCLINIC CLASSES ARE HYPERBOLIC

M. J. PACIFICO, E. R. PUJALS, M. SAMBARINO, J. L. VIEITEZ

ABSTRACT. We shall prove that  $C^1$ -robustly expansive codimension-one homoclinic classes are hyperbolic.

## 1. INTRODUCTION

Let  $M$  be a  $d$ -dimensional manifold and  $\text{Diff}^1(M)$  be the set of  $C^1$  diffeomorphisms  $f$  on  $M$  endowed with the  $C^1$  topology. Let  $p$  be a hyperbolic periodic point of  $f$  and  $H(p)$  be its homoclinic class. The diffeomorphism  $f$  is  $\alpha$ -expansive in  $H(p)$  if  $\text{dist}(f^n(x), f^n(y)) \leq \alpha$  for all  $n \in \mathbb{Z}$ , with  $x, y \in H(p)$ , implies  $x = y$ . It is well known that hyperbolicity implies  $\alpha$ -expansiveness for some  $\alpha > 0$ . But expansiveness alone does not guarantee hyperbolicity, even when one is dealing with a codimension one expansive homoclinic class, as can be seen in [PPV, Section 2]. We note that there are even examples of expansive codimension one homoclinic classes such all of its periodic orbits are hyperbolic that are not hyperbolic.

To see this consider a Smale horse-shoe  $H$  in a plane and  $\Lambda$  a non trivial minimal subset of  $H$ . In a complementary direction multiply  $H$  by a non uniform contraction  $\lambda(w)$ , depending on the distance from  $w$  to  $\Lambda$ , and in such way that  $\lambda(w) = 1$  for  $w \in \Lambda$ . The resulting homoclinic class is expansive, has all periodic points hyperbolic but it is not hyperbolic.

Then we assume that expansiveness holds in a  $C^1$ -neighborhood of the homoclinic class, that is, for all diffeomorphism  $g$   $C^1$  near  $f$  the homoclinic class  $H(p_g)$  of the continuation  $p_g$  of  $p$  is  $\alpha$ -expansive. In [PPV] it was proved that robustly expansive homoclinic classes of a three dimensional manifold are generically hyperbolic. We generalize this result in two ways. First we drop the assumption  $\dim(M) = 3$  and get that robustly expansive codimension-one homoclinic classes have a codimension-one dominated splitting. Second we prove that robustly expansive codimension-one homoclinic classes with a dominated splitting are hyperbolic.

The main step to obtain this result is to prove that center unstable manifolds for all point  $x$  in the homoclinic class  $H(p)$  are true unstable manifolds, that is, center unstable manifolds are dynamically defined. To achieve this it is enough to get this property for periodic points homoclinically related to  $p$ . For this we use a result of Sambarino that controls the behavior of periodic points homoclinically related to  $p$ . Once this is settled, under the hypothesis that the dominated splitting  $E \oplus F$  is not hyperbolic, the fact that center unstable manifolds are dynamically defined allows us to obtain a hyperbolic periodic point  $q$  with arbitrarily large period as near  $H(p)$  as we wish, and with arbitrarily small rate of contraction along the  $E$ -direction. Then we prove that

---

*Date:* January 16, 2006.

M.J.Pacifico and E.R. Pujals were partially supported by CNPq-Brazil/Faperj-Brazil/Pronex Dyn. Systems.

such hyperbolic periodic points are in fact in  $H(p)$ , contradicting uniform  $E$ -contraction in the period for periodic points in  $H(p)$ . As in [Ma2] this implies that the sub-bundle  $F$  is uniformly expanding, and then  $E \oplus F$  is hyperbolic.

## REFERENCES

- [Do] A. DOLD, *Lectures on algebraic topology*, Classics in Mathematics. Springer-Verlag, Berlin., **1995** ISBN: 3-540-58660-1.
- [Fr] J. FRANKS, *Necessary conditions for stability if diffeomorphisms*, Trans. Amer. Math. Soc., **158** (1971), p. 301-308.
- [Ha] S. HAYASHI, *Connecting invariant manifolds and the solution of the  $C^1$ -stability and  $\Omega$ -stability conjecture for flows*, Ann. of Math. (2), Vol. **145** N° **1** (1997), p. 81-137.
- [Li] S.T.LIAO, *On the stability conjecture*, Chinese Ann. Math., **1** (1980), p. 9-30.
- [Ma2] R. MAÑÉ, *A Proof of the  $C^1$  Stability Conjecture*, Inst. Hautes Études Sci. Publ. Math., **66** (1988), p. 161-210.
- [Ma3] R. MAÑÉ, *An ergodic closing lemma*, Annals of Mathematics, **116** (1982), p. 503-540.
- [PM] J. PALIS AND W. DE MELO, *Geometric Theory of Dynamical Systems*, ., **Springer Verlag** 1982..
- [PI1] V. A. PLISS, *Analysis of the necessity of the conditions of Smale and Robbin for structural stability of periodic systems of differential equations*, Diff. Uravnenija, **8** (1972), p. 972-983.
- [PI2] V. A. PLISS, *On a conjecture due to Smale*, Diff. Uravnenija, **8** (1972), p. 268-282.
- [PS] E. PUJALS, M. SAMBARINO, *Homoclinic tangencies and hyperbolicity for surface diffeomorphisms*, Annals of Mathematics, **151** (2000), p. 961-1023.
- [PS2] E. PUJALS, M. SAMBARINO, *On the Dynamics of Dominated Splitting*, Preprint, (2002), p. -.
- [PPV] M. J. PACIFICO, E. R. PUJALS, J. L. VIEITEZ, *Robust expansive homoclinic classes*, Ergodic Theory and Dynamical Systems, **25** (2005), p. 271-300.
- [SV] SAMBARINO, M.; VIEITEZ, J., *On  $C^1$ -persistently Expansive Homoclinic Classes*, Accepted for publication in Discrete and Continuous Dynamical Systems, (2005), p. -
- [1] [We2]WL2 L. WEN, *Heteroclinic Cycles and Homoclinic Closures for Generic Diffeomorphisms*, Preprint, (2002), p. -.